

PLEASE READ ENTIRE LABEL BEFORE USING  
**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

May irritate eyes, nose, throat, and skin. Avoid breathing dust and spray mist. Avoid contact with eyes, nose, throat, skin and clothing. Wash thoroughly after using. Keep away from fire or sparks.

**STATEMENT OF PRACTICAL TREATMENT**


In case of contact, flush skin with plenty of water. For eyes, get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from area being treated. Do not contaminate water by cleaning of spray equipment or disposal of waste. Do not contaminate ornamental ponds.

**STORAGE and DISPOSAL DIRECTIONS**

Never allow Benomyl 50 W (Benlate) to become wet during storage as this may reduce its effectiveness as a fungicide. Keep container closed when not in use. Do not use where staining or damage from water might result. Store in dry place. Do not re-use empty container. Wash thoroughly and dispose of properly.



**ACTIVE INGREDIENTS**  
 Benomyl\*  
 (Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate) 50%  
**INERT INGREDIENTS** 50%  
 U.S. Pat. 3,541,213 & 3,631,176  
 \*Benlate reg. trademark E.I. DuPont de Nemours & Co., Inc.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Side Panel for Additional Precautionary Statements

NET WEIGHT 6 OUNCES

EPA Est. No. 7478-GA-1 EPA Reg. No. 7478-50

DISTRIBUTED BY

CHEM PAK CO  
 Rt 1 Box 114 McRae, GA 31055

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Dip with dry spoon. Do not mix more than amount to be used in a few hours. Keep spray stored.

Apply in sufficient water to obtain thorough coverage, usually 2 to 5 gallons per 1000 sq. ft. For Dollar Spot and Large Brown Patch do NOT water in.

**DOLLAR SPOT** (Small Brown Patch) (*Sclerotinia homocarpa*) Apply at rate of 1 oz. per 1000 sq. ft. at earliest appearance of disease and continue at 10 - 14 day intervals as long as needed.

**LARGE BROWN PATCH** (*Rhizoctonia Solani*) Apply at rate of 2 oz. per 1000 sq. ft. When conditions are unusually favorable for disease, reduce spray intervals from 10 - 14 days to 7 days.

**FUSARIUM PATCH** (Pink Snow Mold or Scald) (*Fusarium nivale*) Apply at rate of 2 oz. per 1000 sq. ft. at earliest appearance of disease and continue at 10 - 14 day intervals as long as needed.

6 level tablespoons equal one ounce.

**Notice:** User assumes all risk of use, storage, or handling of this material not in strict accordance with direction given herewith.