


Date of Draft: 5/14/84

Gustafson  
CAPTAN 300  
Seed Protectant  
Agricultural Fungicide



ACTIVE INGREDIENTS:

Captan (N -- (trichloromethyl)thio-4-cyclohexene-1,  
2-dicarboximide)..... 28.4%  
Related Derivatives..... 1.6%

INERT INGREDIENTS:..... 70.0%

TOTAL..... 100%  
Contains 3 lbs. Captan and related derivatives per U.S. gallon at  
68 F.

KEEP OUT OF REACH OF CHILDREN  
DANGER


- . Causes irreversible eye damage and skin irritation.
- . Harmful if swallowed, inhaled or absorbed through skin.
- . Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.
- . Avoid inhaling spray mist.

See side panel for additional precautionary statements

EPA Est. No. 7501-ID-1  
EPA Reg. No. 7501-8-AA

NET CONTENTS: 5 U.S. Gallons

Gustafson 17400 Dallas North Parkway, Suite 220 --Dallas, Texas  
75252



Read all directions before using and use only as specified on this label.

**DIRECTIONS:** Thoroughly mix the recommended amount of Captan 300 Seed Protectant into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

**IMPORTANT:** Before using -- stir thoroughly to mix contents or use entire container at one time. Do not combine colloidal suspensions, wettable powder or emulsifiable concentrates unless previous experience assures compatibility.

PRODUCT INFORMATION:

Captan 300 Seed Protectant is a flowable concentrate especially formulated for the treatment of seed prior to storage and planting to protect seed from molds and other fungi causing storage loss and to protect seed from seedborne and soil-borne fungi which cause seed decay, damping-off, and seedling blights.

CAPTAN 300  
Approved Application Rate  
Fl. oz. Captan 300 per 100 lbs. of seed

SEED	
Alfalfa, clover, lespedeza, trefoil.....	6-1/2 fl. oz./100 lbs.
Beans (snap, dry, cowpeas).....	2-1/4 fl. oz./100 lbs.
Beets (garden).....	9-1/2 fl. oz./100 lbs.
Cereal grains (for seed and seedling rots - not for bunt smut)	
Barley, rye.....	2 fl. oz./100 lbs.
Oats.....	2-1/3 fl. oz./100 lbs.
Wheat.....	1-1/2 fl. oz./100 lbs.
Cole crops (broccoli, Brussels sprouts, cabbage, cauliflower).....	
.....	1-1/4 fl. oz./100 lbs.
Corn-Field.....	1-1/2 fl. oz./100 lbs.
Sweet.....	3-1/4 fl. oz./100 lbs.
Cotton	
Acid delinted.....	2-1/3 fl. oz./100 lbs.
Reginned, fuzzy.....	3-1/2 fl. oz./100 lbs.
Cucumber, cantaloupe, tomato.....	2-1/3 fl. oz./100 lbs.
Crucifers (collards, kale, mustard, radish, rape, turnips).....	
.....	1-1/4 fl. oz./100 lbs.
Flax.....	3 fl. oz./100 lbs.
Grasses.....	6-1/2 fl. oz./100 lbs.
Peanuts.....	6 fl. oz./100 lbs.
Peas, lentils.....	2 fl. oz./100 lbs.
Pepper (California Wonder).....	2-1/3 fl. oz./100 lbs.
Rice.....	2-3/4 fl. oz./100 lbs.
Sorghum (hulled) for kernel smut.....	3-1/2 fl. oz./100 lbs.
Soybeans.....	2 fl. oz./100 lbs.
Spinach.....	4-3/4 fl. oz./100 lbs.
Squash, pumpkin, watermelon, muskmelon.....	1-2/3 fl. oz./100 lbs.
Sugar Beets	
West.....	4-3/4 fl. oz./100 lbs.
East.....	9-1/2 fl. oz./100 lbs.
Sunflowers.....	4 fl. oz./100 lbs.
Swiss chard.....	9-1/2 fl. oz./100 lbs.

**IMPORTANT**

The Federal Government and most states have seed laws or regulations requiring an appropriate caution statement be affixed to bags containing treated seeds, such as;

**Treated Seed**

Do Not Use for Food, Feed or Oil Purposes.

Mix thoroughly before use or use entire container at one time. Do not combine colloidal suspensions, wetttable powder or emulsifiable concentrates unless previous experience assures compatibility. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

This product is to be used in liquid or slurry treaters similar to Panomatic, Mist-O-Matic or Calkins.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

**USE PRECAUTION**

When using formulations that do not contain dye, to comply with 21CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

