PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE PRECAUTION

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

STORAGE: Store in a cool, dry location. Do not store in direct hot sunlight. After prolonged storage in intense cold, allow product to return to approximately 25 °F. before using.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Gustafson ®

CAPTAN 30-DD

SEED PROTECTANT
FUNGICIDE WITH COLORANT

ACTIVE INGREDIENTS:

*Captan	28.70%
Related Derivatives	0.65%
INERT INGREDIENTS:	70.65%
TOTAL	100.00%

* N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide

Contains 3 lbs. Captan and related derivatives per U.S. gallon at 68°F.

KEEP OUT OF REACH OF CHILDREN
DANGER
STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician. If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention. If on skin: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs.

See side panel for additional precautionary statements.

EPA Est. 7501-ID-1
EPA Reg. No. 7501-9-AA

NET CONTENTS: 5 U.S. Gallons

Gustafson ® is a Reg. TM of Gustafson, Inc.

PRODUCT INFORMATION:

Captan 30-DD Seed Protectant is a flowable concentrate especially formulated for the treatment of small grain, legumes and pulses to protect seed from molds and other fungi causing damage seed and to protect seed from soil borne organisms responsible for fungi which cause seed decay, damping-off, and seedling blights.

CAPTAN 30-DD
Recommended Application Rate
Fl. Oz. Captan 30-DD per 100 lbs. of Seed

Crop	Rate
Alfalfa, clover, lespedeza, vetch	6-1/2 fl. oz./100 lbs.
Beans (snap, dry, limaseed)	3-1/4 fl. oz./100 lbs.
Beets (garden)	6-1/2 fl. oz./100 lbs.
Cereal grains (for seed and seedling rots — not for feed use):	
Barley, rye	2 fl. oz./100 lbs.
Oats	2-1/2 fl. oz./100 lbs.
Wheat	1-1/2 fl. oz./100 lbs.
Cole crops (broccoli, Brussels sprouts, cabbage, cauliflower)	1-1/4 fl. oz./100 lbs.
Corn — Field	1-1/2 fl. oz./100 lbs.
Sweet	3-1/4 fl. oz./100 lbs.
Cotton:	
Acid delinted	3-1/2 fl. oz./100 lbs.
Regrowth, fuzzy	3-1/2 fl. oz./100 lbs.
Cucumber, watermelon, tomato	2-1/2 fl. oz./100 lbs.
Cucurbits (pattypan, kale, mustard, mulch, rape, turnips)	1-1/4 fl. oz./100 lbs.
Flax	3 fl. oz./100 lbs.
Grasses	6-1/2 fl. oz./100 lbs.
Peanuts	6 fl. oz./100 lbs.
Peas, Lentils	3 fl. oz./100 lbs.
Pepper (California Wonder)	2-1/2 fl. oz./100 lbs.
Sorghum (pulses) for forage seed	3-1/2 fl. oz./100 lbs.
Soybeans	2 fl. oz./100 lbs.
Spinach	4-3/4 fl. oz./100 lbs.
Squash, pumpkin, watermelon, muskmelon	1-3/4 fl. oz./100 lbs.
Sugar Beets:	
West	4-3/4 fl. oz./100 lbs.
East	6-1/2 fl. oz./100 lbs.
Sunflowers	4 fl. oz./100 lbs.
Swiss chard	6-1/2 fl. oz./100 lbs.

Read all directions before using and use only as specified on this label.

DIRECTIONS: Thoroughly mix the recommended amount of Captan 30-DD Seed Protectant into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

IMPORTANT: Before using — add thoroughly to mix combine or use only a dispenser at one time. Do not combine other formulations, wettable powder or emulsifiable concentrates unless previous experience assures compatibility.

IMPORTANT: The Federal Government and most states have seed laws or regulations requiring an appropriate caution [illegible] more be affixed to bags containing treated seeds, such as:

Treated Seed
Do Not Use for Food, Feed or Oil Purposes.

When using formulations that do not contain dye, to comply with EPCRR Chapter 1, Section 2.38, all grain seed treated with an organic pesticide must be colored to distinguish seed treated pesticide from untreated seed so as to be used for food or feed for sale. Mix thoroughly before use or use entire contents at one time. Do not combine unused treated seed; store treated seed in a cool place. Do not store in direct sunlight. Follow Seed Handling Instructions.

This product is to be used in fluid or slurry treatment similar to Panoramic, Mini-O-Mitic or Calbrex.

The Federal Food and Drug Administration has not established tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

NOTICE - READ CAREFULLY
LIMITED WARRANTY STATEMENT

Treatment of highly overheated damaged seed, or seed of broken low vigor and poor quality, except for the purpose of certified control of seedling diseases or insect pests, may result in reduced germination and/or reduction of stand and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of retreated seed.

Customer warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach of this limited warranty, Customer shall not be liable for consequential damages.

A quality product manufactured by: Gustafson ® 1400 Preston Road, Suite 400 — Plano, Texas 75093 (214) 985-8872

90(5) WR: [illegible]

06-02-1
[illegible]

BEST AVAILABLE COPY

ACCEPTED
MAR 9 1990
Under the Federal Insect
Fun
c.
to,
EPA [illegible] 7501-9