

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IN CASE OF CONTACT:** Immediately flush eyes with running water for at least 15 minutes. Call a physician. **IN CASE OF SKIN CONTACT:** Wash the affected area with soap and warm water.

**GENERAL USE PRECAUTIONS:**

During commercial exposure in seed treatment wear goggles, clean synthetic rubber gloves, protective clothing and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR part 11.

Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Bury wastes in isolated area away from water supplies.

**IMPORTANT:**

The Federal Government and most states have seed laws or regulations requiring an appropriate caution statement be affixed to bags containing treated seeds, such as:  
Treated Seed

Do Not Use for Food, Feed, or Oil Purposes.

**DIRECTIONS FOR USE  
AND RECOMMENDATIONS:**

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions before using and use only as specified on this label.

**DIRECTIONS:** Thoroughly mix the recommended amount of Captan 400 Seed Protectant into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

**IMPORTANT:** Before using — stir thoroughly to mix contents or use entire container at one time. Do not combine colloidal suspensions, wettable powder or emulsifiable concentrates unless previous experience assures compatibility.

**STORAGE AND DISPOSAL**

1. **PROHIBITIONS**—Do not re-use empty containers. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.
2. **PESTICIDE DISPOSAL**—Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by federal, state or local disposal authorities.
3. **CONTAINER DISPOSAL**—Triple rinse (or equivalent) and dispose of in an approved landfill or bury in a safe place.
4. **GENERAL**—Consult federal, state, or local disposal authorities for approved alternate procedures.
5. **PRODUCT STORAGE**—Store in a cool, dry location. Do not store in direct hot sunlight. After prolonged intense cold storage, allow Captan 400 to return to approximately 25°F before using.

# Gustafson

## CAPTAN 400

**SEED PROTECTANT  
AGRICULTURAL FUNGICIDE**

**ACTIVE INGREDIENT:**

Captan: N[(Trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide. . . . .  
Related Derivatives . . . . .  
**INERT INGREDIENTS:** . . . . .

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

- Causes eye and skin irritation.
- Harmful if swallowed, inhaled or absorbed through skin.
- Do not get in eyes, on skin or on clothing. Wear eye shield and rubber gloves when handling. Wash with water after handling.
- Avoid inhaling spray mist.

**ACCEPTED**  
DEC 29 1980  
Used in the Federal Insecticide, Fungicide, and Rodenticide Act. as a pesticide.  
EPA Reg. No. 7501-26

Contains 4 Pounds Captan Per Gallon  
EPA Est. No. . . . .  
EPA Reg. No. 7501-26 **23** NET CONTENTS

BEST DOCUMENT AVAILABLE

# Gustafson

## CAPTAN 400

SEED PROTECTANT  
AGRICULTURAL FUNGICIDE

**ACTIVE INGREDIENT:**

Captan: N[(Trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide ..... 37.0%  
Related Derivatives ..... 2.1%

**INERT INGREDIENTS:** ..... 60.9%  
..... 100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

- Causes eye and skin irritation.
- Harmful if swallowed, inhaled or absorbed through skin.
- Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.
- Avoid inhaling spray mist.

See side panel for additional precautionary statement.

ACCEPTED  
EPA 29-1980  
Federal Insecticide, Fungicide and Rodenticide Act  
EPA Reg. No. 7501-26

Contains 4 Pounds Captan Per Gallon

EPA Est. No.

EPA Reg. No. 7501-26 **23**

11/80

NET CONTENTS: 5 U.S. Gallons

**PRODUCT INFORMATION:**

Captan 400 Seed Protectant is a flowable concentrate especially formulated for the treatment of seed prior to storage and planting to protect seed from molds and other fungi causing storage loss and to protect seed from seedborne and soil-borne fungi which cause seed decay, damping-off, and seedling blights

Captan 400	
Approved Application Rate Range	
Fl. oz. Captan 400 per 100 lbs. of seed	
<b>SEED</b>	—Range—
Alfalfa, clover, lupine, vetch, trefoil	5 fl. oz.—8 fl. oz./100 lbs.
Beans (snap, dry cowpeas)	2 fl. oz.—3 fl. oz./100 lbs.
Beets (garden)	8 fl. oz.—12 fl. oz./100 lbs.
Cereal grains (for seed and seedling rots-not for bunt smut)	
Barley, rye	2 fl. oz.—3 fl. oz./100 lbs.
Oats	2 fl. oz.—4 fl. oz./100 lbs.
Wheat	1½ fl. oz.—4 fl. oz./100 lbs.
Cole crops (broccoli, Brussels sprout, cabbage, cauliflower)	1 fl. oz.—2 fl. oz./100 lbs.
Corn	
Field	1½ fl. oz.—2-3/8 fl. oz./100 lbs.
Sweet	2 fl. oz.—4 fl. oz./100 lbs.
Cotton	
Acid delinted	2½ fl. oz.—5 fl. oz./100 lbs.
Reginned, fuzzy	3½ fl. oz.—7 fl. oz./100 lbs.
Cucumber, cantaloupe	2½ fl. oz.—3 fl. oz./100 lbs.
Crucifers (Mustard, radish, rape, turnips)	1 fl. oz.—2 fl. oz./100 lbs.
Flax	2 fl. oz.—3-3/4 fl. oz./100 lbs.
Grasses	5 fl. oz.—8-1/4 fl. oz./100 lbs.
Milo	4 fl. oz.—6 fl. oz./100 lbs.
Peanuts	3 fl. oz.—6 fl. oz./100 lbs.
Peas, Lentils	2½ fl. oz./100 lbs.
Peppers	2 fl. oz.—3 fl. oz./100 lbs.
Rice	3 fl. oz.—6 fl. oz./100 lbs.
Sorghum (hulled) For kernel smut	1-3/4 fl. oz.—4½ fl. oz./100 lbs.
Soybeans	1½ fl. oz.—2½ fl. oz./100 lbs.
Spinach	4 fl. oz.—6 fl. oz./100 lbs.
Squash, pumpkin, watermelon, muskmelon	1½ fl. oz.—2 fl. oz./100 lbs.
Sugar Beets	
West	6 fl. oz./100 lbs.
East	12 fl. oz./100 lbs.
Sunflowers	2 fl. oz.—4 fl. oz./100 lbs.
Swiss chard	10 fl. oz.—12 fl. oz./100 lbs.

**NOTICE—READ CAREFULLY  
WARRANTY STATEMENT**

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. In the event of any breach by Gustafson of this limited warranty, Gustafson shall not be liable for consequential damages.

7501-26