FEB 1 1987

Under the Federal Inc.
Fungicide, and Roden:
Fungicide under Roden
Funcion Roden
Funci

`

## PRECAUTIONARY STATEMENTS **HAZARD TO HUMANS** AND DOMESTIC ANIMALS

## CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food.

#### **ENVIRONMENTAL HAZARD**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry location. Do not store in direct, hot sunlight.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

TOPS-2.5D is a Reg. TM of Gustafson, Inc. Gustafson () is a Reg. TM of Gustafson, Inc.

# ®Gustafson ⟨୬⟩ **TOPS** — 2.5D\*

To Sant Company of the

## POTATO SEED - PIECE TREATMEN. **FUNGICIDE DUST**

#### **ACTIVE INGREDIENT:**

Thiophanate-methyl (dimethyl((1,2-phenylene) bis(iminocarbonothiovi)]bis(carbamate))\* .................... 2.5%

INERT INGREDIENTS: ..... 97.5%

TOTAL ......100.0%

\*Also known as dimethyl 4,4'-0-phenylenebis[3-thioallophanate].

## KEEP OUT OF REACH OF CHILDREN



If swallowed, induce vomiting with a mixture of soap or mustard in warm water. Call a physician immediately. If on skin, wash immediately with soap and water.

See Side Panel for additional Precautionary Statements

EPA Reg. No. 7501-32 EPA Est. No. 38091-ID-1 **NET WEIGHT: 50 POUNDS** 

#### **GENERAL INFORMATION**

GUSTAFSON TOPS 2.5D is a fungicide formulated for the dry treatment of potato seed pieces to suppress Fusarium and Rhizoctonia. This pesticide will not control bacterial diseases or other diseases not claimed on this label.

#### **DIRECTIONS FOR USE**

It is a violation of the Federal law to use this product in a manner inconsistent with its labeling. Do not use this product at higher than recommended rate.

For the suppression of potato seed piece decay caused by Fusarium and Rhizoctonia spp., use 1 pound GUSTAFSON TOPS 2.5D per 100 pounds of cut seed pieces. Apply so that the cut seed pieces are thoroughly covered with dust by the use of a convenient container or dust attachment over the belt

Thoroughly clean and sanitize cutting machines, knives, trays, tables, barrels, equipment, trucks and planters before cutting and planting seed pieces. If seed potatoes have not sprouted, subject them to temperatures of 50° to 60°F for about 10 to 15 days before handling. Treat the warmed seed pieces immediately after they are cut. The cut and treated seed pieces should be held for a week or more at cool temperature and high relative humidity; however, they may be planted immediately. This holding period promotes suberization and allows the cut surfaces to heat

Do not use treated seed pieces for food, feed or fodder.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

## NOTICE—READ CAREFULLY LIMITED WARRANTY STATEMENT

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably litter purposes stated on the label only when used in accordance with directions and instructions specified on the label, Gustalson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, quarantee or representation, express or implied, concerning this material. In the event of any breach by Gustalson of this limited warranty, Gustalson shall not be liable for consequential damages.

The state of the state of