ACCEPTED

PRECAUTIONARY **STATEMENTS**

HAZARDS TO HUMANS AND DOMEST & ANIMALS CAUTION

EPA Rag. No.

Harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of nose, throat, eyes and skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Protective clothing (long pants, long-sleeved shirts, gloves, hat and boots) must be worn during mixing and loading. In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Wash hands and face thoroughly with soap and water after use or before eating or smoking. Remove and wash contaminated clothing before reuse

ENVIRONMENTAL HAZARDS

This material is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of federal faw to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

- 1. PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal or by cleaning of equipment. Open dumping is prohibited
- 2. STORAGE: Store in a cool, dry location. Do not store in direct hot sunlight. Keep con tainer closed when not in use
- 3. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be dis posed of on site or at an approved waste disposal facility
- 4. CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state or local authorities

VITAVAX® -MANEB FLOWABLE FUNGICIDE

COMPOSITION

ACTIVE INGREDIENTS: (% by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathiin-3-	
carboxanilide)*	12%
Maneb (Manganese ethylenebisdithiocarbamate)	
Manganese equivalent as metallic 3.51%)	12%
INERT INGREDIENTS:	76%
Total	100%
*U.S. Patents 3,249,499 - 3,393,202 - 3,454,391	

KEEP OUT OF REACH OF CHILDREN CAUTION!

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and see a physician

IF INHALED: See a physician if an abnormal reaction occurs

IF ON SKIN: Wash thoroughly with soap and water

Immediately flush eyes with plenty of water. See a physician IF IN EYES:

See left side panel for Additional Precautionary Statement

EPA Est No 7501-ID-1 EPA Reg. No 7501-39

NET CONTENTS 55 Gallons

* VITAVA " in a Regr TM of UtiliROYA CHEMICAL Div. of Univoyal Inc

Gustafson (O : 5 % Flog TN & GG stafson The

A quality product manufactured for

Gustason 4) 17400 Dallas North Parkway, Suite 220 — Dallas, Texas 75252-4493

istafson 🕪

VITAVAX® -MANEB OWABLE FUNGICIDE

COMPOSITION

INGREDIENTS: (% by weight)	
xin (5,6-dihydro-2-methyl-1,4-oxathiin-3-	
oxanilide)*	12%
o (Manganese ethylenebisdithiocarbamate)	
iganese equivalent as metallic 3.51%)	12%
NGREDIENTS:	76%
· · · · · · · · · · · · · · · · · · ·	100%
atents 3,249,499 - 3,393,202 - 3,454,391	

KEEP OUT OF REACH OF CHILDREN **CAUTION!**

STATEMENT OF PRACTICAL TREATMENT

.OWED: Induce vomiting and see a physician

See a physician if an abnormal reaction occurs

Wash thoroughly with soap and water

Immediately flush eyes with plenty of water. See a physician

See left side panel for Additional Precautionary Statement

No. 7501-ID-1 No. 7501-39

NET CONTENTS

55 Gallons

a RegrITM of UNIROYAL CHEMICAL, Div. of Uniroyal Inc.

is a fleg TN of Gustatson the.

DIRECTIONS FOR USE -T

GUSTAFSON VITAVAX-MANEB Flowable Fungicide is a ready-to-use seed treatment.

RICE

GUSTAFSON VITAVAX-MANEB Flowable Fungicide combines the systemic action of carboxin with the surface action of maneb to control various seed and seedling diseases of rice. It is particularly effective against Rhizoctonia solani and Helminthosporium oryzee.

BARLEY, WHEAT AND OATS ~

In addition, loose smut (Ustilago nude) and covered smut (Ustilago hordel) are controlled on barley, and loose smut (Ustilago tritici) and common bunt (Tilletia carles) are controlled on wheat by GUSTAFSON VITAVAX-MANEB Flowable Fungicide.

Use 4.5-6 fluid ounces of GUSTAFSON VITAVAX-MANEB Flowable Fungicide per 100 pounds of seed. GUSTAFSON VITAVAX-MANEB Flowable Fungicide should be applied as a slurry treatment by machine

SOYBEANS

Use 3 fluid ounces of GUSTAFSON VITAVAX-MANEB Flowable Fungicide per bushel of seed or 6 fluid ounces per 100 pounds of seed. GUSTAFSON VITAVAX-MANEB Flowable Fungicide should be applied as a sturry treatment by machine

USE RESTRICTIONS

Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting. Care must be exercised in the handling of treated seed.

When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180 1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals

COMPATIBILITY

GUSTAFSON VITAVAX MANEB Flowable Fung-cide should not be tank mixed with any pesticide unless adequate physical compatability and phytotoxicity tests have been completed and proved to be satisfactory

SEED LABELING -- "This seed tre-ited with GUSTAFSON VITAVAX-MANEB Flowable Fungicide. Do not use treated seed for food, feed or oil purposes. Add appropriate information for

NOTICE - READ CAREFULLY LIMITED WARRANTY STATEMENT

Gustafson warrants that this product conforms to the chemical description on the label and is real-onably fit for purposes stated on the label only when used in accordance with directions and instructions splicified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach by Gustafson of this limited warranty, Gustafson shall not be liable for consequential damages