GUSTAFSON APRON®-FL SEED TREATMENT

ACCIONO D with COMP LATS in EVA Late, Plant

Fungicide

Version of Act perticide A Reg. No.

FEB 2 4 1983

A seed treatment chemical for control of seed rot and damping off diseases of cotton and certain other crops.

For use only by commercial seed treaters

5 US Gallons Net Weight

Keep Out of Reach of Children.

Warning

See additional precautionary statements on left panel.

EPA File Symbol 7-01-

•, •,

THRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crom injury, ineffectiveness, or other unintended consequences may result to ause of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GUSTAFSON, INC. or the Seller. All such risks shall be assumed by the Buyer.

Gustafson warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes, any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach by Gustafson of this limited warranty, Gustafson shall not be liable for consequential damages.

General Information

GUSTAFSON APRON-FL is a systemic fungicide seed dressing specifically for control of Pythium and Phytophthora spp. For control of other soilborne diseases, such as Rhizoctonia spp. GUSTAFSON APRON-FL should be applied in combination with other registered seed dressing fungicides.

GUSTAFSON APRON-FL is a systemic fungicide having a specific mode of action. GUSTAFSON APRON-FL could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible GUSTAFSON APRON-FL resistant strains of fungi which may occur.

GUSTAFSON APRON-FL may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW DIFECTIONS ON THIS LABER MAY RESELT IN DROP INJURY OR POUR DISEASE CONTROL.

Cotton	For Pythium seed rot and damping off: Apply GUSTAFSON
	APRON-FE as a seed treatment at the rate of .75-1.50
	fluid ounces per 100 lbs. of seed.

For Pithilm damping off and early season Phytophthesis control: Jobbly GCSTH 22% NP 3 N-FL as a seed territo int at the rate of .75-1.80 fluid nances per 100 lbs. of are.1.

Altalia, claser, lestedaza, beans (for for ape), and forage Legimen beans (for forage), soybean may, pelmuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet

beans (for forage).

Southering

	For Pythium damping off and early season Phytophthora control: Apply GUSTAFSON APRON-FL as a seed treatment at the rate of .75-1.50 fluid nunces per 100 lbs. of seed.
Grain Crops	Wheat, barley, rye, oats, sorghum (grain), millet, milo, rice, corn (field corn, sweet corn and popcorn), and buckwheat.
	For Pythium damping off control: Apply GUSTAFSON APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Seed and Pod Vegetables	Black-eyed peas, cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils. For Pythium damping off and early season Phytophthora control: Apply GUSTAFSON APRON-FL as a seed treatment
	at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Ryegrass	For Pythium damping off control on golf courses and home lawns: Apply GUSTAFSON APROM-FE as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Forage Grasses	Grasses grown for hay grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay grazing or silage.
	For Pythium damping off control: Apply GUSTAFSON APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.

Note:

Federal law requires that bags containing treated seeds shall be labeled with the following information: "This seed has been treated with GUSTAFSON APRON-FL fungicide. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs." Use with an EPA-approved dye that imparts an unnatural color to the seed.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
- 2. <u>PESTICIDE DISPOSAL</u> Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- 3. CONTAINER DISPOSAL Triple rinse (or equivalent) and dispose in an inclnerator or landfill approved for pesticide containers, or bury in a safe place.
- 4. GENERAL Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

Precautionary Statements

Hazards to Humans and Domestic Animals

WARTING

Causes eye irritation. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes. Avoid contact with skin or clothing. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse. Do not store near food or feed.

Practical Treatment: In case of contact, flush eyes or skin with plenty of water. If swallowed, call a physician intentately. Do not induce vomiting. If inhaled, move to fresh air; if not breathing, give artificial resultation and get medical attention.

Environmental Marchigo De not apply describe to water. Apply only as specifically the easy of Denot apply when weather interpretation, taken into the mitroated areas. No not continued water by idearing at equipment or disposal of weather.

Physical interests the extens policy of a most consequence the specifical.

APRON' trademark of CIBA-GEIGY for metalaxyl

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (#H-467)		DATE OF ISSUANCE
ASHINGTON, L.C. 21440	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: REGISTRATION	NAME OF PESTICIDE PRODU-	СТ
(Under the Federal Insectivide Function) and Rodentis (de Act. a) ananded)		
AME AND ADDRESS OF REGISTRANT (Include ZIP code)		,
Г	٦	1.1
		· /.
		111 215-2
	t	11 1.
	7	
OTE: Changes in labeling formula differing in substance f ubmitted to and accepted by the Registration Division prior roduct always refer to the above U.S. EPA registration number on the basis of information furnished by the registrant, the	or to use of the label in commerciber.	ce. In any correspondence on the
on the basis of information furnished by the registrant, the he Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is here	oy kegistered/keregistered unda
copy of the labeling accepted in connection with this Re	gistration/Reregistration is ret	urned herewith.
Act is not to be construed as giving the registrant a right to by others.	o exclusive use of the maine of	to at use if a mas been covered
ATTACHMENT IS APPLICABLE		

EPA Form 8570-6 (Rev. 5-76)

PREVIOUS ECITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.