

PM 21
7501-42

D. L. McFadden
Gustafson, Inc.
P. O. Box 660065
Dallas, TX 75266-0065

OCT 25 1995

Dear Mr. McFadden:

Subject: Apron®-FL seed Treatment Fungicide (EPA Reg. No. 7501-42)
Label Amendment
Correspondence Dated May 30, 1995
Correspondence Dated June 26, 1995


The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration or reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Should you have any questions or concerns, please call Kathryn Scanlon on (703) 305-7382.

Sincerely Yours,



Connie B. Welch
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

7505C:K.Scanlon:ks:10/16/95:ApronFL2.LTR

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	K. Scanlon							
DATE ▶	Oct 16, 1995							

Oct 11 1995

Gustafson APRON® -FL

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

7501-42

Seed Treatment Fungicide

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED
ROT AND DAMPING OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENT:

Metalaxyl: N-(2,6-dimethylphenyl)-
N-(methoxyacetyl)alanine methyl ester 28.35%
INERT INGREDIENTS: 71.65%

TOTAL 100.0%

Contains 2.65 pounds Metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Additional Precautionary Statements.

EPA Reg. No. 7501-42

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. May cause eye irritation. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton, and shoes plus socks. Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

If on skin: Wash affected area with soap and warm water.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

PROHIBITIONS—Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

STORAGE—Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL—Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL—Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation.

IMPORTANT: Read the entire **Directions for Use** and the **Limited Warranty Statement** before using this product.

Gustafson



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton.

NOTE: Federal law requires that bags containing treated seeds shall be labeled with the following information. "This seed has been treated with Gustafson APRON-FL fungicide. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs." Use with an EPA approved dye that imparts an unnatural color to the seed.

Gustafson APRON-FL is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, combination of Gustafson Captan and VITAVAX® registered formulations are compatible. Do not use with other seed treatment products unless previous experience assures compatibility. Gustafson APRON-FL may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type seed treatment equipment.

Note: Gustafson APRON-FL is a systemic fungicide having a specific mode of action. Gustafson APRON-FL could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Gustafson APRON-FL resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

Cotton	For <i>Pythium</i> seed rot and damping-off: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Soybeans	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Sorghum	For control of <i>Pythium</i> and systemic downy mildew: Apply Gustafson APRON-FL as a seed treatment at the rate of 1.5-3.0 fluid ounces per 100 lbs. of seed to hybrids with resistance to at least two pathotypes of downy mildew. The 3.0 fl. oz. rate is only recommended for area with a recent severe history of a pathotype of downy mildew that is pathogenic to the hybrid being

3 of 4
treated. Seed treated with the 3.0 fl. oz. rate should be planted and not carried over to the following year. Only high quality sorghum seed (90% germ and above) should be treated with Gustafson APRON-FL.

Sunflowers	For control of systemic downy mildew: Apply Gustafson APRON-FL as a seed treatment at the rate of 3 fluid ounces per 100 lbs. of seed.
Beets, Garden Carrots Sugar Beets	For control of <i>Pythium</i> damping-off: Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Forage Legumes	Alfalfa, clover, lespedeza beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Grain Crops	Wheat, barley, rye, oats, millet, milo, rice, corn (field corn, sweet corn and popcorn), buckwheat and triticale. For <i>Pythium</i> damping-off control: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.
Seed and Pod Vegetables	Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils. For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply Gustafson APRON-FL as a seed treatment at the rate of 3.4 fluid ounces per 100 lbs. of seed.
Cucumbers	For control of <i>Pythium</i> damping-off: Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed. For season long control, apply RIDOMIL® 2EC or 5G in accordance with label directions.
Turf Grasses	For <i>Pythium</i> damping-off control on golf courses home lawns, reclamation and erosion control projects: Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Forage Grasses	Grasses grown for hay grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay grazing or silage. For <i>Pythium</i> damping-off control: Apply Gustafson APRON-FL as a seed treatment at the rate of .75-1.50 fluid ounces per 100 lbs. of seed.

Seed Treatment-Export Use:

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Sunflowers	Apply Gustafson APRON-FL at the rate of 10 fluid ounces (3.3 ounces active) per 100 pounds of seed.
Corn	Apply Gustafson APRON-FL at the rate of 3 to 10 fluid ounces per 100 pounds of seed.
Cucurbit Vegetables	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Fruiting Vegetables (Except Curcubits)	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.

4 of 4

DIRECTIONS FOR USE (continued)

Leafy Vegetables (Except Brassica)	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Brassica (Cole) Leafy Vegetables	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Root and Tuber Vegetables	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed. NOTE: Do not apply APRON-FL to potatoes.
Onions	Apply Gustafson APRON-FL as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.

Seed treated in accordance with the export use pattern must be labeled as follows:

**FOR EXPORT ONLY-
NOT FOR DOMESTIC SALES OR USE.**

DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

Gustafson APRON-FL is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. Gustafson APRON-FL may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

Corn, Field Sorghum, Grain Wheat Barley	To aid in control of seed decay and damping-off caused by <i>Pythium</i> : Apply Gustafson APRON-FL as a commercial seed treatment at the rate 0.10 fluid ounces per 100 lbs. seed. Apply only in combination with EPA registered rates of Gustafson Flowable Captan products.
Rice	To aid in control of seed decay and damping-off caused by <i>Pythium</i> : Apply APRON-FL as a commercial seed treatment at the rate 0.10 to 0.15 fluid ounces per 100 lbs. of seed. Apply only in combination with EPA registered rates of VITAVAX-200 Flowable or Gustafson 42-S Thiram or RTU-VITAVAX-Thiram Fungicides. Recommended use rates in combination with Gustafson APRON-FL are: VITAVAX-200 FF: 3.0 to 4.0 fluid ounces per 100 lbs. of seed. Gustafson 42-S Thiram: 2.0 to 3.3 fluid ounces per 100 lbs. of seed. RTU-VITAVAX-Thiram: 5.0 to 6.8 fluid ounces per 100 lbs. of seed.

NOTICE - READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson, Inc., warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON, INC. NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson, Inc. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product manufactured by:
Gustafson, Inc.
1400 Preston Road, Suite 400
Plano, Texas 75093
(214) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson, Inc.
APRON® and RIDOMIL® are trademarks of CIBA-GEIGY for Metalaxyl
VITAVAX is a Reg. TM of Uniroyal Chemical Company, Inc.