UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Dennis McFadden Gustafson, LLC 1400 Preston Road, Suite 400 Plano, TX 75093

Subject: Your July 12, 1999 application for label amendment EPA Reg. No. 7501-42 Gustafson Allegiance - FL Seed Treatment Fungicide

Dear Mr. McFadden,

The label amendment above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable providing you make the following revision.

On page three, under "Directions for Use at Reduced Rates in Combination with Other Fungicides, the entry for "Peanuts" must be asterisked and the following statement must be added to the peanut use directions "* Not registered for use in California."

| CONCURRENCES | | | | | | | | |
|--------------|---------|--|--|--|--|-----------------------------------------|--|--------------|
| SYMBOL, | 7505C | | | | | | | · |
| SURNAME | Wwange) | | | | | , ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, | | |
| DATE | 3/31/00 | | | | | | | |
| | | | | | | | | 11 50 5 6001 |

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

OFFICIAL FILE COPY

A stamped copy of the accepted label is enclosed for your records. You must submit one copy of the final printed label to the Agency prior to distribution of this label in channels of trade. If there are questions concerning this matter, contact Dr. Tom Ellwanger at (703) 308-9352.

Sincerely,

Mary L. Waller

Product Manager (21)

Fungicide Branch

Registration Division (7505C)

Mary J. Waller

ACCEPTED with COMMENTS In EPA Letter Dated

Gustafson ALLEGIANCE™-FL

APR 1 1 2000

Seed Treatment Fungicide

Under the Federal Insetticide, Fundicate, and Rodenticide Act

as amendon for the pesticide SEED TREATMENT CHEMICAL FOR CONTROL OF SEED TSOI-42. ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENT:

 Metalaxyl:N-(2,6-dimethylphenyl) 28.35%

 N-(methoxyacetyl)alanine methyl ester
 71.65%

 TOTAL
 100.0%

Contains 2.65 pounds Metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

See Additional Precautionary Statements.

EPA Reg. No. 7501-42 EPA Est. No.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT(PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants, socks and chemical-resistant footwear and chemical-resistant gloves (such as nitrile, butyl, neoprene, and/or barrier laminate). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE AND DISPOSAL

PROHIBITIONS—Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

STORAGE—Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL—Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposed facility.

CONTAINER DISPOSAL—Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty Statement before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: if the seed is treated with the product and the treated seed is soilinjected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, socks and chemical resistant footwear, chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton and protective eyewear.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before being sold. Federal law requires that bags containing treated seeds shall be labeled with the following information. "This seed has been treated with ALLEGIANCE-FL fungicide. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs."

ALLEGIANCE-FL is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora spp.* For control of other soilborne diseases, combination of Gustafson Captan and VITAVAX® registered formulations are compatible. Do not use with other seed treatment products unless previous experience assures compatibility.

ALLEGIANCE-FL may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

Note: ALLEGIANCE-FL is a systemic fungicide having a specific mode of action. ALLEGIANCE-FL could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible ALLEGIANCE-FL resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

| Carrots* | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cotton | For Pythium seed not and damping-off: Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. |
| Soybeans | For Pythium damping-off and early season Phytophthora control: Apply ALLEGIANCE-FL as a seed treatment at the rate of 0.75-1.5 ft. oz. per 100 lbs. of seed. |
| Spinach* | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed. Do not apply ALLEGIANCE-FL to spinach which is destined to be grown in the greenhouse. Containers or bags containing ALLEGIANCE-FL treated spinach seed must be labeled with this restriction. *Not registered for use in California. |
| Peanuts* | For Pythium seed rot and damping-off control: Apply ALLEGIANCE-FL as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California. |
| Sorghum | For control of <i>Pythium</i> damping-off apply 0.375-0.75 fl. oz. of ALLEGIANCE-FL per 100 lbs. of seed applied as a seed treatment. For control of systemic downy mildew, apply 0.75 fl. oz. ALLEGIANCE-FL per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5 fl. oz. of ALLEGIANCE-FL per 100 lbs. of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with ALLEGIANCE-FL. |
| Sunflowers | For control of systemic downy mildew; Apply ALLE-GIANCE-FL as a seed treatment at the rate of 1.5-3.0 fl. oz. per 100 lbs. of seed. |
| Beets, Garden Sugar Beets | For control of <i>Pythium</i> damping-off: Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. |
| Forage Legumes | Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). For Pythium damping-off and early season Phytophthora control: Apply ALLEGIANCE-FL as a seed treatment at the rate of 0.75-1.5 ft. oz. per 100 lbs. of seed. |
| Grain Crops | Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale. |
| | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75 ft. oz. per 100 lbs. of seed. For systemic downy mildew control of sweet com: Apply ALLEGIANCE-FL as a seed treatment at the rate of 1.5 ft. oz. per 100 lbs. of seed. |
| Rice* | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs, of seed. If using ALLEGIANCE-FL treated rice do not use Ridomil for control of Pythium in water-seeded rice. *Not registered for use in California. |
| Seed and Pod Vegetables | Black-eyed peas, chickpeas (Garbanzns), co wpeas, dill, edible soybeans, field beans, field peas, alarden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils. For Pythium damping-off control:Apply ALLEGIANCF-FL as a seed treatment at the rate of 0.75 fl. oz, per 100 lbs. |

| | of seed. For early season <i>Phytophthora</i> control: ApplyALLEGIANCE-FL as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply ALLEGIANCE-FL as a seed treatment at the rate of 1.5 fl. oz. per 100 lb. of seed. |
|----------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cucumbers* | For control of <i>Pythium</i> damping-off. Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California. |
| Turf Grasses | For Pythium damping-off control on golf courses, home lawns, reclamation and erosion control projects: Apply ALLEGIANCE-FL as a seed treatment at the rate of 1.5 ft. oz. per 100 lbs. of seed. |
| Forage Grasses | Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage. For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. |
| | SEED TREATMENT-EXPORT USE: |
| For treatmen | t of the following seed for export only to countries |
| requiring | treatment with metalaxyl before importation. |
| Brassica (Cole) Leafy Vegetables | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 lbs. of seed. |
| Canola | For Pythium damping-off control: Apply Allegiance-FL as a seed treatment at the rate of 0.50 - 1.70 fl. oz. per 100 lbs. of seed. (32 - 110 ml per 100 kg of seed) |
| Com | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed. |
| Cucurbit Vegetables | For Pythium damping-off control: Apply ALLE- GIANCE-FL as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed. |
| Fruiting Vegetables | For Pythium damping-off control: Apply ALLE- GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. |
| Leafy Vegetables | For Pythium damping-off control: Apply ALLE- GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 fbs. of seed. |
| Onions | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. |
| Millet | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed. |
| Peas | For Pythium damping-off and early season Phytophthora control: Apply ALLEGIANCE-FL as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed. |
| Root and | For Pythium damping-off control: Apply ALLE- |
| Tuber Vegetables | GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. |
| | Do not apply ALLEGIANCE-FL to potatoes. |
| Sorghum | For Pythium damping-off control: Apply ALLE-GIANCE-FL as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed. |
| Sunflowers | For systemic downy mildew control: Apply ALLE-GIANCE-FL as a seed treatment at 8.65-10.35 fl. oz. per 100 lbs. of seed. |

Seed treated in accordance with the export use pattern must be labeled as follows:

FOR EXPORT ONLY-NOT FOR DOMESTIC SALES OR USE.

DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

ALLEGIANCE-FL is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. ALLEGIANCE-FL may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

| Com, Field Wheat,Oats, Barley, Rye | To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL as a commercial seed treatment at the rate 0.10-0.375 fl. oz. per 100 lbs.seed. Apply only in combination with EPA registered rates of Gustafson broad-spectrum seed treatment fungicides. | |
|----------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Soybeans, Field beans, Kidney beans, Navy beans, Pinto beans, Lentils Peas | To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL as a commercial seed treatment at the rate 0.10-0.375 fl. oz. per 100 lbs. of seed. Apply in combination with EPA registered rates of broad-spectrum seed treatment fungicides. | |
| Sunflowers | To aid in the control of seed decay and damp ing-off caused by <i>Pythium</i> : Apply ALLE-GIANCE-FL as a commerical seed treatment at the rate 0.10-0.375 fl. oz. per 100 lbs. of seed. Apply in combination with EPA registered rates of broad spectrum seed treatment fungicides. | |
| Peanuts | To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLE-GIANCE-FL as a commercial seed treatmen at the rate 0.10-0.375 fl. oz. per 100 lbs. of seed. Apply in combination with EPA registered rates of braod spectrum seed treatmen fungicides. | |
| Rice* | To aid in control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL as commercial seed treatment at the rate of 0.10 to 0.375 ft. oz. per 100 lbs. of seed. *Not registered for use in California | |

NOTICE - READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing are made to guarantee germination of carry-over seed. As its sole express warranty, Gustafson LLC variants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE, EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLCSHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW GUSTAFSON LLOSA

NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product manufactured by: Gustafson LLC 1400 Preston Road, Suite 400 Plano, Texas 75093 Questions? (972)985-8877

RIDOMIL® is a trademark of NOVARTIS Crop Protection Inc. for mefenoxam ALLEGIANCE is a TM of Gustafson LLC Gustafson (logo) is a Reg. TM of Gustafson LLC

42(S)-008230