

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

7501-RO

DATE OF ISSUANCE

MAY 30 1991

TERM OF ISSUANCE

Conditional

NAME OF PESTICIDE PRODUCT

RTM 5-15 Seed Protectant

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Gustafson, Inc.
P.O. Box 220065
Dallas, TX 75222

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data or other material required for registration, reregistration of your product under FIFRA section 3(c)(7) and FIFRA section 3(c)(7)(A) when the Agency requires all registrants of similar products to submit such data.

2. Add the phrase "EPA Registration No. 7501-RO" to your label before you release the product for shipment.

3. Submit five (5) copies of your final printed labeling to us as you release the product for shipment. Refer to the 4-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 3(c). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Larley
Product Manager (FIFRA)
Fungicide-Pesticide Branch
Registration Division (TS-767)

Enclosures

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

MAY 30 1991

2524

ACCEPTED
with COMMENTS
EPA Letter Dated:

Gustafson

MAY 30 1991

RTU[®]-BAYTAN[®]-Thiram Fungicide

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

7501-80

For Use Only By Commercial Seed Treaters

ACTIVE INGREDIENTS:

Beta-(4-Chlorophenoxy)-alpha-(1,1-dimethylethyl)	
-1H-1,2,4-triazole-1-ethanol	5.0%
Thiram (Tetramethylthiuram disulfide)	15.3%
INERT INGREDIENTS	79.1%
TOTAL	100.0%

Contains 0.43 pounds triadimenol (BAYTAN) and 1.33 pounds Thiram per U. S. gallon.
U. S. Patent No. 3,952,002

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

EPA Reg. No. 7501-
Net Contents: 2 1/2 Gallons

EPA Est No. 7501-ID-1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid inhaling of spray mist or dust. Do not use or store near food or feed. This product may cause allergic reactions.

During commercial exposure in seed treatment and loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety and Health Administration (formerly the U. S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under provisions of 30 CFR, Part 11. Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work, wash all clothing with soap and water before reusing. Consumption of alcoholic beverages increases the toxic effects of Thiram

(Cont. Next Column)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Administer water freely and induce vomiting by giving one dose (1/2 oz. of 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately. **IF ON SKIN:** Wash skin immediately with soap and warm water. If irritation persists, get medical attention. **IF EYES ARE CONTAMINATED:** Wash with plenty of water for at least 15 minutes. If irritation persists, get medical attention. **IF INHALED:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

SYMPTOMS OF POISONING: No specific symptoms. Acute poisoning is accompanied by general depression and illness. Consumption of alcoholic beverages with exposure to Thiram may cause flushing, breathing difficulty, nausea, vomiting and low blood pressure. In case of poisoning, call a physician immediately. Have patient lie down and keep quiet. **NOTE TO PHYSICIAN:** No specific antidote is available. Treat symptomatically. In the presence of consumption of alcoholic beverages, Thiram responds similar to Disulfiram.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water when disposing of equipment washwaters. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Gustafson RTU-BAYTAN-Thiram Fungicide combines the systemic action of BAYTAN with the surface action of Thiram to control or suppress various seed, seedling and soilborne diseases of wheat, barley, oats, rye and corn. Systemic activity of BAYTAN will provide early season control of infections by powdery mildew and rusts.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place.

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SEED TREATMENT: Gustafson RTU-BAYTAN-Thiram Fungicide is a ready-to-use seed treatment. Application should be made using standard slurry or mist-type commercial seed treatment equipment. Uniform application of seed is necessary to insure seed safety and best disease control. Seed should be sound and well cured prior to treatment.

Application of rates less than 6 fluid ounces of Gustafson RTU-BAYTAN Thiram Fungicide per 100 pounds of seed must be diluted with sufficient water to give a minimum applied treatment rate of 6 fluid ounces of slurry/cwt. Rates of 6 to 9 ounces of product may be applied without further dilution.

NOTE: To meet U. S. Federal Seed Act requirements, all seed treated with RTU-BAYTAN-Thiram Fungicide should be labeled: "Treated with beta-(4-chlorophenoxy)-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol (BAYTAN) and Tetramethylthiuram disulfide (Thiram). Do not use treated seed for feed, food or oil purposes. Do not graze green forage within 40 days of planting."

RESTRICTIONS

For use only through commercial seed treating equipment which are closed application systems. Do not use treated seed for feed, food or oil purposes.

RECOMMENDED APPLICATIONS

CROP	DISEASE	FLUID OUNCES RTU-BAYTAN-Thiram Fungicide	REMARKS
Wheat Barley Oats Rye	Seed and Soilborne Diseases:	4-1/2 to 9 fluid ounces per 100 pounds of seed	Refer to the Seed Treatment section on this label for the recommended method of treating seed. Apply the 4-1/2 ounce rate for control of smuts, and the 9 ounce rate for control or suppression of the other diseases. *Suppression or some control of these diseases provided. Green forage may be grazed 40 days after seeding.
	Wheat:		
	Scab		
	Stinking smut		
	Flag smut		
	Loose smut		
	Glume blotch		
	Foot rot*		
	Take-all*		
	Pythium rot*		
Barley:			
Scab			
Covered smut			
Loose smut			
Net blotch*			
Typhula rot*			
Foot rot*			
Pythium rot*			
Oats:			
Scab			
Loose smut			
Leaf spot*			
Pythium rot*			
Rye:			
Scab			
Stripe smut			
Pythium rot*			

(Cont. Next Page)

40/4

CROP	DISEASE	FLUID OUNCES RTU-BAYTAN-Thiram Fungicide	REMARKS
Wheat Barley Oats Rye (Cont.)	Foliar Diseases: Wheat: Powdery mildew Leaf rust Stripe rust Stem rust Speckled leaf blotch Barley: Barley rust Stripe rust Stem rust Powdery mildew Leaf blotch Oats: Crown rust Stripe rust Stem rust Powdery mildew Rye: Rye rust Stem rust Stripe rust Powdery mildew	9 fluid ounces per 100 pounds of seed	Refer to the Seed Treatment section on this label for the recommended method of treating seed. Use of the recommended rate will control the noted early season diseases. Green forage may be grazed 40 days after seeding.

**NOTICE - READ CAREFULLY
LIMITED WARRANTY STATEMENT**

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach of this limited warranty, Gustafson shall not be liable for consequential damages.

A quality product manufactured by:
GUSTAFSON, INC.
 1400 Preston Road
 Suite 400
 Plano, Texas 75093
 (214) 985-8877

Gustafson (logo) and RTU are Reg TM's of Gustafson, Inc.
 BAYTAN is a Reg. TM of Bayer AG, Germany.