

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

ENVIRONMENTAL HAZARDS

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

Gustafson

VITAVAX®-PCNB

FLOWABLE FUNGICIDE

ACTIVE INGREDIENTS

Carboxin (6,6-dihydro-2H-pyridin-4(1H)-one) 17%

Carboxin (dry) 17%

INERT INGREDIENTS 66%

100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

ESSENTIAL OILS
FISH OIL
FISH MEAL
FISH WASTE

See left side panel for additional precautionary statement.

DIRECTIONS FOR USE

Gustafson Vitavax-PCNB Flowable Fungicide is a ready to use seed treatment. Gustafson Vitavax-PCNB Flowable Fungicide combines the systemic action of carboxin with the surface action of PCNB to control various seed and seedling diseases. It is particularly effective against Rhizoctonia solani.

BARLEY, WHEAT AND OATS
Use 3.4 fluid ounces of Gustafson Vitavax-PCNB Flowable Fungicide per 100 pounds of seed. Gustafson Vitavax-PCNB Flowable Fungicide should be applied as a slurry treatment by machine. Use the higher rate under conditions of expected heavy disease pressure.

In addition, loose smut (rusty spot) and covered smut (rusty spot) are controlled on barley and rice smut (rusty spot) and common bunt (wheat) are controlled on wheat by Gustafson Vitavax-PCNB Flowable Fungicide.

COTTON
Use 5 to 7 fluid ounces of Gustafson Vitavax-PCNB Flowable Fungicide per 100 pounds of seed as a slurry treatment by machine. Use the higher rate under conditions of expected heavy disease pressure. Do not graze or feed livestock on hay grown from treated seed.

SOYBEANS
Use 2 fluid ounces of Gustafson Vitavax-PCNB Flowable Fungicide per bushel of seed or 4 fluid ounces per 100 pounds of seed. Gustafson Vitavax-PCNB Flowable Fungicide should be applied as a slurry treatment by machine.

USE RESTRICTIONS
Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated seed for 30 days after planting. Care must be exercised in the handling of treated seed. Do not store treated seed in a place where seed treated with this product (except by the manufacturer) must be stored with an EPA approved dye (e.g., 40 CFR 160.1000) which identifies the treated seed to help prevent the inadvertent use of treated seed as food or feed for animals.

COMPATIBILITY
Gustafson Vitavax-PCNB Flowable Fungicide should not be tank mixed with any pesticide unless the pesticide manufacturer's directions have been completed and approved by the manufacturer.

SEED LABELING — This seed treated with Gustafson Vitavax-PCNB Flowable Fungicide should be labeled as such for all feed or food purposes. Add appropriate information for any other than a treatment.

**NOTICE — READ CAREFULLY
LIMITED WARRANTY STATEMENT**

This limited warranty does not cover damaged seed or seed of known low vigor and poor quality. The purchase of this product for the control of existing disease or insect pests may result in a reduction of yield and/or reduction of seed and seedling vigor. Treat and conduct germinal tests in a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

This limited warranty does not apply to seed which does not conform to the chemical description on the label and which is not used in accordance with the directions on the label. Gustafson, Inc., neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY. Guarantee or representation expressed or implied concerning this material in the event of any breach of this limited warranty. Gustafson shall not be liable for consequential damages.

ACCEPTED

NOV 24 1987

the Federal Insecticide
Fungicide and Rodenticide Act,
for the pesticide
registered under
EPA Reg. No. 7501-87