

7501-91

09/19/2000

508206 1/3

-1-

ACCEPTED
 SEP 19 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7501-91

BAYTAN® 30

Flowable Fungicide

For Use Only By Commercial Seed Treaters

ACTIVE INGREDIENT (% by weight)

Triadimenol**.....	30%
INERT INGREDIENTS:.....	70%
TOTAL:.....	100%

**Contains 2.65 pounds Triadimenol per gallon
U.S. Patent No. 3,952,002

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements.

EPA Reg. No. 7501-91 EPA Est.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Do not breathe vapors. Applicators must wear long pants, long-sleeved shirts and chemical resistant gloves. Do not get in eyes, on skin or on clothing. Do not contaminate feed or food. Wash hands, arms and face thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

(Cont. Next Column)

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

- Call a poison control center or doctor for treatment advice.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover treated seeds that are visible on soil surface, especially in end rows and turn areas. Clean up spilled treated seeds to reduce exposure to wildlife.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place.

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Gustafson



RECOMMENDED APPLICATIONS

CROP	DISEASE	FLUID OUNCES BAYTAN 30 Flowable Fungicide	REMARKS
Wheat Barley Oats Rye	Seed and Soilborne Diseases:	0.75 to 1.5 fluid ounces per 100 pounds of seed	Refer to the Seed Treatment section on this label for the recommended method of treating seed. Apply the 0.75-ounce rate for control of smuts, and the 1.5-ounce rate for control or suppression of the other diseases. *Suppression or some control of these diseases provided. Green forage may be grazed 40 days after seeding.
	Wheat: Stinking smut Flag smut Loose smut Glume blotch Foot rot* Take-all* Barley: Covered smut Loose smut Net blotch* Typhula rot* Foot rot* Oats: Loose smut Rye: Stripe smut		
	Foliar Diseases:	1.5 fluid ounces per 100 pounds of seed	Refer to the Seed Treatment section on this label for the recommended method of treating seed. Use of the recommended rate will control the noted early season diseases. Green forage may be grazed 40 days after seeding.
	Wheat: Powdery mildew Leaf rust Stripe rust Barley: Barley rust Stripe rust Powdery mildew Oats: Crown rust Stripe rust Powdery mildew Rye: Rye rust Stripe rust Powdery mildew		
Corn	Head smut	3 fluid ounces per 100 pounds of seed	Refer to the Seed Treatment section on this label for the recommended method of treating seed. Green forage may be grazed 45 days after seeding.
Cotton (Acid Delinted)	<i>Rhizoctonia</i> Seedling Disease <i>Thielaviopsis basicola</i> (Black root rot)	0.25 to 0.75* fl. oz. per 100 lbs. of seed 1.0 to 3.0 fl. oz. per 100 lbs. of seed	* May be used in combination with VITAVAX-PCNB or Vitavax-30C for added protection against <i>Rhizoctonia</i> seed and seedling disease. For protection against seedling disease caused by <i>Thielaviopsis basicola</i> . Higher rates are only for areas with a severe history of disease.

BAYTAN 30 Fungicide controls or suppresses certain designated seed and soilborne diseases of wheat, barley, oats and rye. Systemic activity of BAYTAN 30 Fungicide will provide early season control of infections by powdery mildew and rusts.

SEED TREATMENT: BAYTAN 30 Fungicide may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 16 fl. ozs. of water per 100 lbs. of seed. Add dye** to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

**21 CFR Chapter I, Section 2.25 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored with an EPA -approved dye, such as one of the dyes listed on 40 CFR Section 180.1001(c) and (d), to prevent their subsequent inadvertent use as a food for man or feed for animals.

NOTE: In the absence of significant disease pressure, seed treated with BAYTAN may emerge slightly slower than non-treated. This effect usually disappears two to three days after initial emergence. For best results, do not plant seed at depths greater than 1-1/2 inches.

SEED LABELING

To meet U. S. Federal Seed Act requirements, all seed treated with BAYTAN Seed Treatment Fungicide should be labeled: "Treated with Triadimenol (BAYTAN)."

RESTRICTIONS

For use only through commercial seed treating equipment which are closed application systems. Do not use treated seed for feed, food or oil purposes.

NOTICE - READ CAREFULLY

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product manufactured by:
GUSTAFSON LLC
1400 Preston Road
Suite 400
Plano, Texas 75093
Questions?(972) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson LLC
BAYTAN is a Reg. TM of Bayer AG, Germany.
Vitavax is a Reg. TM of Uniroyal Chemical Co., Inc.

91(S)-005070