



PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid inhaling of dust. Do not use or store near food or feed.

Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work; wash all clothing with soap and water before reusing.

Consumption of alcoholic beverages increase the toxic effects of thiram.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands.

STORAGE AND DISPOSAL

1. STORAGE - Store in a cool place. Keep container tightly closed to prevent evaporation. Stir before using.
2. PROHIBITIONS - Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
3. PESTICIDE DISPOSAL - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
4. CONTAINER DISPOSAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Gustafson VITAVAX Pour-On Flowable Fungicide is a ready-to-use hopper box seed treatment, combining the systemic action of carboxin with the surface action of Thiram to control a broad spectrum of seed and seedling diseases of soybeans. It is particularly effective against seed borne diseases that affect germination and emergence; *Penicillium*, *Aspergillus*, *Diaporthe*, *Phomopsis*, and *Rhizoctonia* sp. VITAVAX Pour-On is applied directly to seed in the hopper box and is safe to peat and granular soybean inoculum.

### SOYBEANS

#### SEED TREATMENT:

Apply 5 fl. oz. of VITAVAX Pour-On Flowable per bushel of seed or 10 fl. oz. per 100 lbs. of seed. Mix well prior to use.

VITAVAX Pour-On Flowable is designed to provide rapid, uniform treatment of soybean seed in the hopper box of unit and drill type planter. Partially fill hopper box with a premeasured amount of seed. Apply the proper amount of product evenly over the surface of the seed. Mix with a stick or paddle until a uniform color is imparted to all seed. Repeat procedure until hopper box is filled.

#### SEED TREATMENT AND INOCULUM STICKER:

VITAVAX Pour-On Flowable is safe to soybean inoculum and will perform as a STICKER for peat or granular inoculum.

Partially fill hopper box with premeasured amount of seed. Apply VITAVAX Pour-On Flowable evenly over the surface of the seed using the appropriate rate per quantity of seed. Mix with a stick or paddle until a uniform color is imparted to all seed. Apply proper amount of inoculum evenly over treated seed, mix thoroughly with stick or paddle. Repeat procedure until hopper box is filled.

#### USE PRECAUTIONS:

1. DO NOT MIX WITH BARE HANDS.
2. Seeding rate should be checked during planting.
3. If seeding operation is interrupted for several hours, stir seed vigorously prior to continuing seeding. Keep hopper box lids closed to prevent dust from entering the hopper box.
4. Do not store seed treated with VITAVAX Pour-On Flowable. Leftover treated seed should be double sown or buried away from water supplies.
5. Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on vines grown from treated seed.