

# ANCHOR®

## FLOWABLE FUNGICIDE

A SEED TREATMENT CHEMICAL FOR CONTROL AND SUPPRESSION OF SEED ROT AND DAMPING OFF DISEASES OF CERTAIN CROPS.

For Use Only By Commercial Seed Treaters

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

**STORAGE:** Do not use or store near heat or open flame.  
**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL

Do not apply this product through any type of irrigation system.

**Note:** ANCHOR Flowable Fungicide is a systemic fungicide having a specific mode of action. ANCHOR Flowable Fungicide could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible ANCHOR Flowable Fungicide resistant strains of fungi which may occur.

ANCHOR Flowable Fungicide is a systemic fungicide seed dressing specifically for control of *Pythium* and suppression of *Phytophthora* spp. For control of other soilborne diseases, such as *Rhizoctonia* spp., ANCHOR Flowable Fungicide should be applied in combination with other registered seed dressing fungicides. ANCHOR Flowable Fungicide may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment. Use the higher rate when conditions of heavy disease pressure are expected.

**NOTE:** Federal Law requires that bags contains treated seeds shall be labeled with the following information. "This seed has been treated with Oxadixyl Fungicide. Do not use treated seed for feed, food or oil purposes." Store away from feeds and food stuffs.

### DIRECTIONS FOR USE (CONT.)

<b>Fruiting Vegetables:</b> Cantaloupes, cucumbers, eggplant, gourds, melons, peppers (including chili, sweet, bell and pimentos), pumpkins, summer squash, tomatoes, watermelons and winter squash	For <i>Pythium</i> damping off: Apply ANCHOR as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
<b>Leafy Vegetables:</b> Broccoli, brussels sprouts, cabbage, cauliflower, celeriac, collards, kale, kohlrabi, lettuce, mustard, parsley, rape, rhubarb and spinach	For <i>Pythium</i> damping off: Apply ANCHOR as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
Soybeans, peas	For <i>Pythium</i> damping off; and early season <i>Phytophthora</i> control; Apply ANCHOR as a seed treatment at the rate of 1.50-2.25 fluid ounces per 100 lbs. of seed.
Cotton (acid delinted, delinted mechanically)	For <i>Pythium</i> seed rot and damping off; Apply ANCHOR as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 lbs. of seed.
Sunflower	For <i>Pythium</i> seed rot and damping off; Apply ANCHOR as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 lbs. of seed. For suppression of early season systemic Downy Mildew, apply at the rate of 3.0 fluid ounces per 100 lbs. of seed.
<b>Root and Tuber Vegetables:</b> Beets, carrots, parsnip, radish, rutabagas, sugar beets and turnip.	For control of <i>Pythium</i> damping off. Apply ANCHOR as a seed treatment at the rate of 1.50 fluid ounces per 100 lbs. of seed.
<b>Grain Crops</b> (except wheat): Barley, buckwheat, corn (field, sweet and popcorn), millet, oats, rye, sorghum and triticale.	For control of <i>Pythium</i> damping off; Apply ANCHOR as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 lbs. of seed.
<b>Turf Grasses:</b>	For <i>Pythium</i> damping off on golf courses and home lawns; Apply ANCHOR as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 lbs. of seed

Forage Grasses:

Forage Legumes:

Alfalfa, clover, lespedeza, lupines, trefoil, vetch and velvet beans.

### USE RESTRICTIONS

Do not use treated seed for food, if treated with this product must be cool (e.g., those listed in 40 CFR 180.10 color) to the seed.

### NOTICE - READ CONDITIONS OF SALE AND STATEMENT

Treatment of highly mechanically damaged, low vigor and poor quality, except for control of existing disease or insect pests, is not a guarantee of germination and/or reduction of seed and seed germination tests on a small portion of total seed lot to a selected chemical conditions beyond the control of Gustafson, Inc. to guarantee germination of carry-over seed. As its sole express warranty, Gustafson, Inc. product conforms to the chemical specifications reasonably fit for purposes stated or indicated with directions and instructions subject to the inherent risks set forth BY APPLICABLE LAW, IN THE EVENT OF LIMITED WARRANTY, GUSTAFSON, INC. IS NOT LIABLE FOR CONSEQUENTIAL DAMAGES. GUSTAFSON, INC. AUTHORIZES ANY OF ITS AGENTS TO MAKE ANY WARRANTY OF FITNESS FOR USE OR REPRESENTATION CONCERNING THIS MATERIAL. Buyer/End User must handle, use and store this product in accordance with instructions and use directions contained on the product label. Buyer/User purchases this product under the terms of Sale and Limited Warranty, which may be varied only by a written agreement authorized representative of Gustafson, Inc. acceptable, return all products to the manufacturer for a full refund.

A quality product manufactured by Gustafson, Inc.  
 1400 Preston Road, Suite 400  
 Plano, Texas 75093

214-985-8877

JUN 23 1994

### ACTIVE INGREDIENT:

Oxadixyl (2-methoxy-N-(2-oxo-1,3-oxazolidin-3-yl)acet-2,6-xylylidide) ..... 29.5%  
 INERT INGREDIENTS ..... 69.0%  
 Total ..... 100.0%

Contains 2.66 pounds oxadixyl per gallon

EPA Reg. No. 7501-97

EPA Est. 7501-ID-1

NET CONTENTS: 2-1/2 U.S. Gallons

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Give one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. **IF IN EYES:** Flush with plenty of water. Call a physician if irritation persists.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label. Cover or incorporate spilled treated seeds

Gustafson (Logo) and ANCHOR are Reg. TM's of Gustafson, Inc.

**Gustafson** 