# ACCEPTE Gustafson ().

FEB | 0 1989

Under the Federal Insecticide, Fungicide, and Rodenucide Act. as amended, for the pesticide segistered under SCI-98

# Reldan® 2%

**DUST INSECTICIDE** 

#### ACTIVE INGREDIENTS:

6

ACTIVE MONICOLENS.	
Chlorpyrifos-methyl [0.0-dimethyl-0-(3,5,6-	
trichloro-2-pyridyl)phosphorothicate]	2%
INERT INGREDIENTS	
This product contains the toxic inert ingredier	t Meth
ylene Chloride.	

#### KEEP OUT OF REACH OF CHILDREN

#### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

- If Swallowed: Do not induce vomiting. Call a physician immediately.
- If Inhaled: Remove victim to fresh air. Apply respiration, if indicated, Call a physician.
- If On Skin: Remove contaminated clothing and immediately wash with soap and water.
- If In Eyes: Flush eyes with plenty of water and call a physician.
- NOTE TO PHYSICIAN: Chlorpyrifos-methyl is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection is the preferable antidote. Oximes such as 2-PAM may or may not be therapeutic, but it is recommended they not be used in place of atropine.

SEE ADDITIONAL PRECAUTIONARY STATE-MENTS

EPA Reg. No. 7501-98 U.S. Patent No. 3,244,586 Est. No.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

CAUSES EYE IRRITATION--MAY CAUSE ALLERGIC SKIN REACTION HARMFUL IF SWALLOWED

Do not get in eyes, on skin, or on clothing. Remove contaminated clothing. Wash thoroughly after handling. Wear goggles or face shield and rubber Cent. Next Column

gloves whenever eye or skin exposure is likely to occur and wash before reuse. Avoid breathing dust. Keep container closed. Handle dust in a ventilated area.

#### **ENVIRONMENTAL HAZARDS**

RELDAN Insecticide is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not discharge directly or indirectly into surface waters. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry location. Do not store in direct, hot sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORED GRAIN: For protection of barley, cats, rice sorghum and wheat against injury from stored grain weevils, moths, beetles and mealworms including granary weevil, rice weevil, red flour beetle, confused flour beetle, saw-toother grain beetle, main meal moth. Angoumois grain moth and suppression of lesser grain borer. Apply to the grain at the rate of 15 lbs. of product per 1000 bu, as the grain is being loaded or turned into final storage, if grain is insect infested, furnigate to course existing populations and apply RELDAN for residual protection. To protect stored grains from attack by Indian meal moth, apply to the surface of clean or infested grain at the rate of 7 lbs. per 1000 sq. ft. of grain surface area. Apply evenly. Repeat if necessary.

# HOW TO APPLY TO SMALL GRAINS

Apply Gustafson RELDAN 2% Dust insecticide when grain is in the truck or wagon prior to binning. To determine the amount in the truck or wagon, multiply the length by width by depth (all in feet) and multiply this total by 0.8 to get the number of bushels. Measure out the correct amount of Gustafson RELDAN2% Dust insecticide (10 lbs. per 1000 bu.) and spread evenly over the surface of the grain in the truck or wagon. Cut into the grain with a shovel. Coating will become mixed with grain as it is dumped from the truck or wagon into the hopper. Further mixing will occur when the grain is augered or blown into the bin. Mixing is completed as the coating and grain roll down the cone shaped pile in the bin.

\*Furnigate with Phosphine or Methyl Bromide according to manu-facturers label recommendations.

## **USE PRECAUTIONS:**

Avoid application at extremely dusty sites.

Check equipment calibration regularly to insure proper application rates.

When drying grain, apply RELDAN after the grain has cooled.

### NOTICE--READ CAREFULLY LIMITED WARRANTY STATEMENT

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or Implied, concerning this material. In the event of any breach of this limited warranty, Gustafson shall not be liable for consequential damages.

A quality product manufactured by: Gustafson, Inc. 1400 Preston Road, Suite 400 Plano, Texas 75075 (214) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson, Inc. RELDAN is a Reg. TM of The Dow Chemical Company

L131S-8/88