



BEST AVAILABLE COPY

**SEED TREATMENT
INSECTICIDE-FUNGICIDE**

Active Ingredients:

Carboxin (5,6-dihydro-2-methyl-2-oxo-1,2,4-triazin-3-carboxamide)	14%
Diazinon (0,6-dimethyl-2-isopropyl-6-methyl-4-pyrimidinyl phosphorothioate)	15%
Lindane (gamma isomer of benzene hexachloride)	25%
Inert Ingredients:	46%

100%

Contains Graphite

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See additional precautionary statements

EPA Reg. No. 7501-112
EPA Est. 420564L-1

**PRECAUTIONARY STATEMENTS
Hazards to Humans and
Domestic Animals
CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on clothing. Applicators must wear long-sleeved shirt, long pants, gloves, and a disposable paper dust mask which covers at least one-third of the face. Wash hands thoroughly after using and before eating or smoking. This product should be applied in a well-ventilated area. Avoid contamination of feed and foodstuffs. Do not store with food. Do not store with feed or forage to be fed to dairy animals or to animals being finished for slaughter.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions before using and use only as specified on this label.

APR 12 1961

7501-112

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry place. Do not allow to freeze or become overheated.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Germate Plus VITAVAX-Diazinon-Lindane Seed Treatment is a ready to use dust base product especially formulated for seed treatment at planting time. This product is formulated to help lubricate the planter box and assure a smooth even flow to the furrow. It contains a systemic fungicide VITAVAX for protection of seeds from seed and seedling diseases, Diazinon for protection against seed corn beetles and seed corn maggots, and Lindane for protection against wire worms. Germate Plus VITAVAX-Diazinon-Lindane Seed Treatment may be used on seed previously treated with Captan or Thiram. Do not use on seed previously treated with any insecticide other than a stored grain protectant. Do not use on foundation seed stock.

CORN, SOYBEANS, BEANS

Planter Box Application: Use Germate Plus VITAVAX-Diazinon-Lindane Seed Treatment at the rate of 1.5 pounds (one pound per 50 pounds of seed) (2.0 lb/BU). Apply as a planter box treatment mixing thoroughly with the seed before planting. For best results the following procedure is suggested: Fill the planter box only 1/2 full of seed, and run the required amount of product and mix thoroughly in the box. Then add the remainder of seed to the planter box and add the rest of the required amount of product to the seed, again mixing thoroughly.

Good results are also obtained by premixing seed with the product in a suitable container, then pouring treated seed into the planter box.

Note: Do not use on foundation or seed stock. Do not use on seed already treated with an insecticide other than a stored grain protectant. Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds sufficient to reduce the rate of delivery through a planter, 1 ounce of graphite per bushel of seed should be added as a lubricant. Adjust the planter to give the required rate of seed delivery with the treated seed. Check planter periodically to verify the proper planting rate is being delivered.

Clean-up for Electric Eye Population Monitors: At the end of each day of use, clean out the electric eye monitors with brushes provided by the planter manufacturer, according to the directions.

USE RESTRICTIONS

Treated seed must not be used for animal feed, for food or animal feed, or processed for oil. Care must be exercised in the handling of treated seed. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess seeds beyond planting time.

Read entire label use and disposal directions. Precautionary statements and directions also on the outside of the bag and the label report.

NOTICE - READ CAREFULLY LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach of this limited warranty, Gustafson shall not be liable for consequential damages.

High quality product manufactured by
GUSTAFSON, INC.
1400 Preston Road
Suite 400
Piano, Texas 75075
(214) 965-8877

Gustafson Reg. is a Reg. TM of Gustafson, Inc.
GERMATE is a TM of Gustafson, Inc.
VITAVAX is a Reg. TM of Agrivox Chemical Company, Inc.

112511-8882

Applicators must wear long sleeved shirt, long pants, gloves, and a disposable dust mask covering at least one third of face. This product must be applied in a well ventilated area.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water.

BEST AVAILABLE COPY