
Gustafson ⟨୬⟩

APRON® TERRACLOR®



4-19-90

DUST SEED TREATMENT FUNGICIDE

WITH GRAPHITE

ACTIVE INGREGIENTS:

Pentachloronitrobenzene (PCNB)	 	25 00%
Metalaxyl	 	. 625%
INERT INGREDIENTS:	 	68.75%

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements

EPA Reg No 7501-126

EPA Est 42056-IL-1

Net Contents:

Pounds

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Protective clothing including dust mask, gloves, long-sleeved shirt, long parts, socks and shoes should be worn during mixing/loading procedures.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Cull a physician of Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If In eyes: Flush eyes with plenty of vater. Call a physician if Irritation persists. If on akin: Washwiti plenty of soap and water. Call iii physician if Irritation occurs. If Inhaled: Move to fresh ak. Administer CPR if not breathing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and equatic organisms. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Cover or incorporate treated seeds. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Do not store in direct sunlight. Protect from extreme temperatures.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse contain:

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for quidance

CONTAINER DISPOSAL: Completely empty container Intuapplication equipment. Then offer for recycling or reconditioning, or puncture and dispose of In a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTE: Metalaxyl is a systemic fungicide having a specific mode of action. Metalaxyl could be subject to devulopment of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance or ways to control any possible metalaxyl resistant strains of fungi which may occur.

RECOMMENDED APPLICATIONS

APRON-TERRACLOR will control damping off, seed and seeding rot diseases caused by *Pythium spp.* and *Rhizoctonia solani*. APRON-TERRACLOR will also provide early season control of *Phytophthora* in soybeans.—

PLANTER BOX TREATMENT

Apply at the specified rate and premix with the seed directly in the planter box at planting. A non-oil lubricant has also been added to help seeds flow more easily and evenly from the planter box. Use the higher rate of application in fields with a history of severe disease pressure.

SEED TREATER APPLICATION

APRON-TERRACLOR may be applied at planting time, utilizing the Gustafson CF-80* treater to maximize seed coverage. (*Calibrate the Gustafson CF-80 as per the manufacturer's instructions). Do not use the CF-80 for applications to peanuts.

CROP5	RATE	DISEASE	
Cotton	8 oz./cwt.	Pythium and Rhizoctoria Seediing Disease Complex, Common Smut or Bunt (Wheat), Covered Smut (Barley)	
Peanuts	4 to 8 oz./cwt.		
Barley, Rice	2 to 4 oz./bu.		
Beans, Corn, Wheat	2 oz./bu.		
Soybeans, Peas	4 oz./bu	Pythium and Rhizoctonia Seedling Disease Complex, Early Season Phytophthora	

NOTE: One heaping scoop (enclosed) is equivalent to approximately 4 avoirdupois ounces of thes product.

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. All seed treated with this product (except by the planter box method) must be colored with an EPA approved dye (e.g., those listed in 40 CFR 180.1001) which imparts an unnatural color to the seed.

NOTICE-READ CAREFULLY LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling viljor. Treat and conduct germination tests on a small portion of seer before committing the total seed lot to a selected chemical treatment pue to seed quality conditions beyond the control of Gustatson, Inc., no claims are made to guarantee germination of car.y- over seed.

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. In the event of any breach of this limited warranty, Gustafson shall not be liable for consequential damages.

A quality product manufactured by: Gustafson, Inc. 1400 Preston Road, Suite 400 Plano, Texas 75075 (214) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson, Inc. Terracior is a Reg. TM of Univoyal Chemical Company, Inc. Apron® is a Trademark of Ciba Geigy for Metalaxyl

126(S)-093040

BEST AVAILABLE COPY

13/