

VITAFLO™-280**Flowable Fungicide****ACCEPTED**

1/6/2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7501-133

Active Ingredients:

Carboxin (5,6-dihydro-2-methyl-N-phenyl-1,4-oxathiin -3-carboxamide)	14.9%
Thiram (Tetramethylthiuram disulfide)	13.2%
Inert Ingredients:	71.9%
Total:	100.0%

EPA Reg. No. 7501-133 EPA Est.

NET CONTENTS:

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes:

•Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

•Call a poison control center or doctor for treatment advice.

If swallowed:

•Immediately call a poison control center or doctor.

•Do not induce vomiting unless told to do so by a poison control center or doctor.

•Do not give any liquid to the person.

•Do not give anything to an unconscious person.

If inhaled:

•Move person to fresh air.

•If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

•Call a poison control center or doctor for further treatment advice.

If on skin or clothing:

•Take off contaminated clothing.

•Rinse skin immediately with plenty of water for 15 - 20 minutes.

•Call a poison control center or doctor for treatment advice.

Note to Physician: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

See Additional Precautionary Statements.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid inhaling of spray mist. Do not use or store near food or feed. Consumption of alcoholic beverages increases the toxic effects of thiram. During commercial exposure in seed treatment and loading operations, wear goggles, clean

rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety and Health Administration (formerly the U. S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under provisions of 30 CFR, Part 11. Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work; wash all clothing with soap and water before reusing.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift or runoff may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for use on agricultural establishments in hopper-box, planter-box, slurry-box, or other seed treatment applications at or immediately before planting.

STORAGE AND DISPOSAL

Do not store VITAFLO-280 in direct sunlight. Do not store VITAFLO-280 at temperatures above 95° F. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

VITAFLO-280 is a combination of a systemic fungicide (VITAVAX®) and a contact fungicide (Thiram) providing plant protection against seed and soilborne seedling diseases including damping-off and seed decay. In addition, VITAFLO-280 provides protection against the following cereal grain diseases: Barley: covered smut (*Ustilago hordei*), true loose smut (*U. nuda*) suppression of net blotch (*Helminthosporium teres*). Canola: *Rhizoctonia solani*, *Pythium* seed and seedling decay, *Phoma* (black leg) and *Alternaria* blight. Wheat:

Gustafson 

loose smut (*U. tritica*), common bunt (*Tilletia caries*). Oats: loose smut (*U. kolleria*), covered smut (*U. avenae*). VITAFLO-280 has no vapor action. Thorough seed coverage is required. Use the higher rate under conditions of heavy expected disease pressure. VITAFLO-280 can be diluted with water for use in slurry treaters. Dilute VITAFLO-280 as required to apply the specified rate of undiluted VITAFLO-280. Keep in constant agitation and use within 24 hours of diluting. Treater calibration should be rechecked if the VITAFLO-280 temperature falls below -4° F. Undiluted VITAFLO-280 can be used at temperatures down to -22° F. Centrifugal pumps are not recommended for pumping VITAFLO-280. Use closed loading system.

TREATMENT RATE FOR UNDILUTED VITAFLO-280

CROP	fl. oz. per cwt. seed
Barley, Wheat (The 3.5 fl. oz. rate will give partial control of true loose smut in wheat and barley. Use the 5 fl. oz. rate for control of true loose smut).	3.5 - 5.0
Canola, rapeseed	8.8
Oats	5.0
Triticale	3.0
Corn (field & sweet)	4.5
Soybeans, Beans	4.0

CORN HEAD SMUT: For protection against seedborne head smut (*Sphacelotheca reiliana*). Treat corn at 8.5 - 11.5 fl. oz. of UNDILUTED VITAFLO-280 per cwt. of seed.

GENERAL NOTE: Avoid planting corn under cold, wet conditions. Stress conditions during the first few weeks after planting may increase the incidence of head smut. Will not control soil-borne head smut.

USE RESTRICTIONS:

- VITAFLO-280 will not freeze solid even in extreme winter temperatures.
- For cleaning lines and reservoirs, flush with water. Rinse water should be used for dilutions of VITAFLO-280.
- Seeding rate should be checked during planting.
- Stir well before using.
- Prior to each use, stir contents for approximately 2 minutes with the internal hand operated agitator.
- Do not use treated seed for food, feed or oil purposes.
- Do not graze or feed livestock on treated areas for six weeks after planting. Care must be exercised in the handling of treated seed. Do not graze or feed livestock on vines grown from treated seed.
- Rape greens grown and harvested from VITAFLO-280 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from VITAFLO-280 treated seed is only for industrial uses and can not be used for edible oil or any other human / feed consumption.

USE PRECAUTION

When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.25 all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180.1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals.

COMPATIBILITY

VITAFLO-280 Flowable Fungicide should not be tank mixed with any pesticide unless adequate physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.

SEED LABELING

"This seed treated with carboxin and thiram (VITAFLO-280 Flowable Fungicide) at the manufacturer's recommended rate. Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated area for six weeks after planting

grain crops. Do not graze or feed livestock on vines grown from treated bean or soybean seed." Add appropriate information for any other chemical treatments.

**NOTICE - READ CAREFULLY
CONDITIONS OF SALE AND LIMITED WARRANTY
STATEMENT**

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label. Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product manufactured by:
Gustafson LLC
1400 Preston Road, Suite 400
Plano, Texas 75093
Questions?(972) 985-8877

VITAFLO is a trademark of Crompton Corporation, Inc.
GUSTAFSON (logo) is a registered trademark of Gustafson LLC,
Plano, Texas