ALLEGIANCETM-TERRACLOR®-VITAVAX® **Fungicide**

ACTIVE INGREGIENTS:

Carboxin: (5,6-Dihydro-2-methyl-N-phenyl-1,	
4-oxathiin-3-carboxamide)	15.00%
Pentachloronitrobenzene (PCNB)	15.00%
Metalaxyl	3.12%
INERT INGREDIENTS:	
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements.

EPA Reg. No. 7501-145

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. PERSONAL PROTECTIVE EQUIPMENT(PPE):

Applicators and other handlers (other than mixers and loaders) must wear: long-sleeved shirt and long pants, waterproof gloves and shoes plus socks. Mixers and Loaders must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention. IF INHALED: Remove victim to fresh air. If not ACCEPTED

JAN 30 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act. as amended, for the pesticide *agistered unday

breathing, give artificial respiration, preferably mouth-tomouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Cover or incorporate treated seeds. Do not contaminate water when disposing of equipment washwaters.

Groundwater Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Do not store in direct sunlight. Protect from extreme temperatures.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for

CONTAINER DISPOSAL: Completely empty container into application equipment. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE....

It is a violation of Federal law to use this productin a manner inconsistent with its labeling.

Do not apply this product in away that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not use treated seed for food, feed or oil purposes.

On corn, barley, oats and wheat, do not graze or feed livestock on treated areas for six weeks after planting. On cotton, peanuts, beans and soybeans, do not graze or feed livestock on hay grown from treated seed. Do not hog down peanuts grown from treated seed.

USE RESTRICTIONS

All seed treated with this product (except by the planter box method) must be colored with an EPA approved dye (e.g., those listed in 40 CFR 180.1001) which imparts an unnatural color to the seed.

NOTICE - READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON, INC. NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FIT-NESS OR MERCHANTABILITY, GUARANTY OR REPRESEN-TATION, EXPRESS OR IMPLIED, CONCERNING THIS MATE-RIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson, Inc. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontainments, notification, and emergency assistance. gency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves and shoes plus socks and protective

NOTE: Metalaxyl is a systemic fungicide having a specific mode of action. Metalaxyl could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Gustafson, Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance or ways to control any possible metalaxyl resistant strains of fungi which may occur.

RECOMMENDED APPLICATIONS

PREVAILALLEGIANCE-TERRACLOR-VITAVAX Fungicide will protect against damping-off, seed and seeding rot diseases caused by Pythium spp. and Rhizocionia solani.

PLANTER BOX TREATMENT

Apply at the specified rate and premix with the seed directly in the planter box at planting. A non-oil lubricant has also been added to help seeds flow more easily and evenly from the planter box. Use the higher rate of application in fields with a history of severe disease pressure

SEED TREATER APPLICATION

PREVAIL may be applied at planting time, utilizing an on-farm mechanical treater* to maximize seed coverage. (*Calibrate as per the manufacturer's instructions.) On peanuts, use only mechanical treaters that will not physically damage peanut seed.

CROPS	RATE	DISEASE	
Cotton	4 to 8 oz./cwt.	For protection against Pythium and Rhizoctonia Seedling Disease Com-	
Peanuts	2 to 4 oz./cwt.		
Barley, Oats, Rice	1 to 2 oz./bu.	plex, Loose Smut and Common Bunt or Stinking Smut (Wheat), Loose and Covered Smut (Barley), Oat Smut	
Beans	3 to 4 oz./cwt		
Corn , Wheat	1.5 oz./bu.		
Soybeans	2 oz./bu.	For protection against Pythium and Rhizoctonia Seedling Disease Complex	

A quality product manufactured fof: Gustafson, Inc. 1400 Preston Road, Suite 400 Plano, Texas 75093

PREVAIL is a Reg. TM of Gustafson, Inc.
TERRACLOR and VITAVAX are Reg. TM's of Uniroyal Chemical Company, Inc. ALLEGIANCE is a TM of Gustafson, Inc.

145(S) 793201