

7501-155

9/2/2001

Gaucho 480 • Master label

1/4

ACCEPTED
0885 10/15/01 Tests and
stored grain

OCT 2 2001

**GAUCHO® 480
FLOWABLE**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7501-155

ACTIVE INGREDIENT:
imidacloprid, 1-[6-chloro-3-pyridinyl)methyl]-N-nitro
-2-imidazolidinimine.....40.7%
OTHER INGREDIENTS:.....59.3%
Total:.....100.0%
Contains 4 lbs. active per gallon (480 grams per liter) @ 20°C.
EPA Reg. No. 7501-155 EPA Est.

**KEEP OUT OF REACH OF CHILDREN
CAUTION****PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

Harmful if swallowed, absorbed through skin or inhaled.
Avoid breathing vapor or spray mist. Wash thoroughly with
soap and water after handling. Remove contaminated cloth-
ing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product
are listed below. If you want more options, follow the instruc-
tions for Category C on an EPA chemical resistance category
selection chart.

Applicators and other handlers must wear: Long-sleeved
shirt and long pants, socks and shoes and chemical-resis-
tant gloves (such as nitrile, butyl, neoprene, barrier laminate,
polyvinyl chloride or Viton). Follow manufacturer's instruc-
tions for cleaning/maintaining PPE. If no such instructions for
washables, use detergent and hot water. Keep and wash
PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing
gum, using tobacco or using the toilet. Users should remove
clothing immediately if pesticide gets inside. Then wash
thoroughly and put on clean clothing. Users should remove
PPE immediately after handling this product. Wash the out-
side of gloves before removing. As soon as possible, wash
thoroughly and change clothing.

FIRST AID

Have the product container or label with you when calling a
poison control center or doctor, or going for treatment.

If swallowed:

-Call a poison control center or doctor immediately for
treatment advice.

-Have person sip a glass of water if able to swallow.

-Do not induce vomiting unless told to do so by a poison
control center or doctor.

-Do not give anything by mouth to an unconscious person.

If on skin or clothing:

-Take off contaminated clothing.

-Rinse skin immediately with plenty of water for 15 - 20
minutes.

-Call a poison control center or doctor for treatment advice.

If inhaled:

-Move person to fresh air.

-If person is not breathing, call 911 or an ambulance, then give
artificial respiration, preferably mouth-to-mouth if possible.

-Call a poison control center or doctor for further treatment
advice.

If in eyes:

-Hold eye open and rinse slowly and gently with water for 15
- 20 minutes. Remove contact lenses, if present, after the first
5 minutes, then continue rinsing eye.

-Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates.
For terrestrial uses, do not apply directly to water, or to areas
where surface water is present or to intertidal areas below the
mean high water mark. Do not contaminate water when
disposing of equipment washwaters. Cover or incorporate
spilled treated seeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

For use in commercial seed treaters only, with the exception of
application to canola, cotton (delinted seed), field corn, sor-
ghum, wheat and barley, which may be made either by commer-
cial seed treatment or as an end-use seed treatment on agricul-
tural establishments at, or immediately before, planting. This
product is to be used in liquid or slurry treaters.

Do not apply this product in a way that will contact workers or other
persons, either directly or through drift. Only protected handlers
may be in the area during application. For any requirements
specific to your State or Tribe, consult the agency responsible for
pesticide regulation.

Mix thoroughly before use or use entire container at one time. All
tank mixes should be pre-tested to determine physical compat-
ibility between formulations. Observe all cautions and limita-
tions on labeling of all products used in mixtures.

STORAGE AND DISPOSAL

STORAGE: Store in a cool place. Do not store in direct sunlight.
Protect from freezing temperatures.

PROHIBITIONS: Do not contaminate water, food or feed by
storage, disposal, or by cleaning of equipment. Do not reuse
empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL: Wastes resulting from the use of this
product may be disposed of on site or at an approved waste
disposal facility.

CONTAINER DISPOSAL: Contact Gustafson or your chemical
supplier for container return instructions. If container is not
returned in accordance with instructions, triple rinse (or
equivalent). Then offer for recycling or reconditioning, or
puncture and dispose of in a sanitary landfill, or incineration,
or, if allowed by state and local authorities, by burning. If
burned, stay out of smoke.

Gustafson

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks, protective eyewear.

Stored Seed Protection:

Applied at labeled rates equal to or above 1.0 fl. oz. per hundredweight, GAUCHO 480 Flowable will provide protection to seed against injury from the following insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). It is recommended that seed with existing populations of stored grain pests be fumigated prior to treating and bagging seed.

For Early Season Protection Against Certain Sucking Insects: GAUCHO 480 Flowable will aid in the protection of seeds and seedlings against injury by certain early season insects.

Canola, Rapeseed

To provide early season protection of seedlings against injury by aphids, flea beetles and wireworms apply as a commercial seed treatment at 12.8 to 32 fl. oz. per hundredweight of seed. In areas where flea beetle and foliar insects are in high numbers, the higher application rate is recommended. Please consult your local agricultural office for pest patterns, history, and forecasts to assist in determining the appropriate rate for your region.

For suppression of Lygus bugs, including *Lygus spp.*, in their second generation and cabbage Seedpod Weevil larvae, including *Ceutorhynchus assimilis*, apply 19.2 to 32 fl. oz. per hundredweight of seed.

Canola, Rapeseed: For End-Use Application At Agricultural Establishments:

Apply using an HCBT or a Batch Treater. Shake GAUCHO 480 Flowable thoroughly before use. Apply 6.4 to 16 fl. oz. of GAUCHO 480 Flowable per 50 pound bag. GAUCHO 480 Flowable may be diluted with an approved Gustafson fungicide mixture for extended disease protection. Treat one-half of seed with one-half of slurry mix. Add the balance of the seed and apply balance of slurry. Allow mixing until seed is thoroughly covered.

Wheat, Barley:

To provide early season protection of seedlings against injury by aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly and Wireworms, apply 1.0 to 3.0 fluid ounces per hundredweight of seed prior to planting as a slurry treatment. Low use rate for "wireworm-only" protection: GAUCHO 480 applied at the 0.16 - 0.32 fl. oz. per hundredweight of seed offers suppression of wireworm activity on seed and young seedlings. Ensure thorough coverage. Use the higher rate to provide increased length of protection and from heavy

insect pressure and from wireworms, and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors. Do not graze or feed livestock on treated areas for 45 days after planting. **Wheat, Barley: For End-Use Application At Agricultural Establishments:**

Apply using a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.), Gustafson Air Pressure System (GAP) or other on-farm seed treating equipment to deliver accurate rates of GAUCHO 480 Flowable to achieve optimum product performance. Apply 1.0 - 3.0 fl. oz. per hundredweight of seed. GAUCHO 480 Flowable should be combined with a Gustafson fungicide product for seed and seedling protection against fungal pathogens, as well as insect pests. Depending on fungicide formulation used dilution with water may be necessary for optimum coverage. GAUCHO 480 Flowable may also be applied on-farm as an over-treatment to seed pretreated with a fungicide. In this case, dilution is necessary. Do not graze or feed livestock on treated areas for 45 days after planting.

Sorghum:

To provide early season protection of seedlings against injury by Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Fire ants and Wireworms, apply 8 fluid ounces per hundredweight of seed prior to planting as a slurry treatment. Ensure thorough coverage. Do not graze or feed livestock on treated areas for 45 days after planting. **Sorghum: For End-Use Application At Agricultural Establishments:**

Apply using an HCBT or an Eight-Bag Batch Treater. Shake GAUCHO 480 Flowable thoroughly before use. Dilute 4 fluid ounces of GAUCHO 480 Flowable with water. The final slurry rate should be adjusted to apply a rate of 8 - 10 fluid ounces of diluted slurry per 50 pound bag of seed. Treat one-half of seed with one-half of slurry mix. Add the balance of the seed and apply balance of slurry. Allow mixing until seed is thoroughly covered. Apply 0.75 oz. of dry TALC per 50 pound bag of seed following the GAUCHO 480 Flowable application and allow it to distribute evenly on the seed. Do not graze or feed livestock on treated areas for 45 days after planting.

Sugar Beets:

To provide early season protection of seedlings against injury by Whitefly, Aphids, Leafhoppers (including to reduce potential for spread of Curly top and Yellow mosaic virus due to aphid and leafhopper vectors), Root aphid, Thrips and Wireworms, apply as a commercial seed treatment at 3.0 to 6.3 fluid ounces of GAUCHO 480 Flowable in or on a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed - approximately 1 kilogram by weight). Apply in a film coat directly to raw seed (100,000 seed or approximately 1 kilogram by weight) at a rate of 3.0 fluid ounces per unit of seed. If rates exceed 3.0 fluid ounces per unit, seed must be pelleted.

Cotton (Delinted Seed Only):

To provide early season protection of seedlings against injury by early season thrips and aphids, apply 6 to 8 fluid ounces per hundredweight of seed prior to planting as a slurry treatment so as to ensure thorough coverage.

Cotton (Delinted Seed Only): For End-Use Application At Agricultural Establishments:

Apply using an HCBT or an Eight-Bag Batch Treater. Shake GAUCHO 480 Flowable thoroughly before use. Dilute 3 to 4 fluid ounces of GAUCHO 480 Flowable with water or a ready to use fungicide mixture, such as Gustafson RTU®-VITAVAX®-Thiram or ALLEGIANCE™-FL. The final slurry rate should be adjusted to apply a rate of eight to ten fluid ounces of diluted slurry per 50 pound bag of seed. Treat one-half of seed with one-half of slurry mix. Add the balance of the seed and apply balance of slurry. Allow mixing until seed is thoroughly covered.

(Continued next page)

Field Corn:

For the protection of corn plants from the corn insect pests listed below. Apply as a seed treatment at the recommended rates listed.

PEST	GAUCHO 480	
	MG. AI / KERNEL	FL. OZ. / 80,000 UNIT OF SEED
Corn root worm (including Northern, Western, Southern and Mexican) ¹ Flea beetle Chinch bug Southern green stinkbug White grub Seed corn maggot Thrips Wireworm Corn leaf aphid Imported fire ant Southern corn leaf aphid Billbug ² Grape colaspis ² Black cutworm ³	1.34	7.6

PEST	GAUCHO 480	
	MG. AI / KERNEL	FL. OZ. / 80,000 UNIT OF SEED
Flea beetle Chinch bug Seed corn maggot Thrips Wireworm Corn leaf aphid Imported fire ant Grape colaspis ² White grub	0.6	3.4

PEST	GAUCHO 480	
	MG. AI / KERNEL	FL. OZ. / 80,000 UNIT OF SEED
Seed corn maggot (seed protection only) Wireworm (seed protection only) Flea beetle (through 1 leaf stage) Imported fire ant White grub ⁴	0.16	0.91

¹In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn rootworm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations.

²Reduces early season feeding damage.

³Will reduce feeding damage caused by leaf feeding black cutworms that are 1/2 inch or less in length.

⁴Reduces feeding damage during emergence and seedling stages.

(Continued next column)

Field Corn: For End-Use Application At Agricultural Establishments:

Apply using an HCBT or an Elght-Bag Batch Treater. Shake GAUCHO 480 Flowable thoroughly before use. Dilute GAUCHO 480 Flowable with water and/or an approved Gustafson fungicide mixture. The final slurry rate should be adjusted to apply a rate of 8 - 10 fl. oz. of diluted slurry per 50 pound bag of seed. Treat one-half of seed with one-half of slurry mix. Allow mixing until seed is thoroughly covered. Apply 0.75 oz. of dry TALC per 50 pound bag of seed following the GAUCHO 480 Flowable application and allow it to distribute evenly on the seed.

Sweet Corn:

For the protection of sweet corn plants from the insect pests listed below. Apply as a seed treatment at the recommended rates listed.

PEST	GAUCHO 480 FL. OZ. / CWT. OF SEED
Flea beetle Early season corn leaf aphid Seed corn maggot Wireworm	8
Imported fire ant Early season corn leaf aphid Seed corn maggot Wireworm	4 - 8
Imported fire ant Seed corn maggot (seed protection) Wireworm (seed protection)	2 - 4
Wireworm (seed protection)	1 - 2

The final slurry rate should be adjusted to apply a rate of 16 - 20 fl. oz. of dilute solution per hundredweight of seed with commercial application equipment. (Not registered for use in California) Succulent shelled beans (including runner bean, snap bean, wax bean, asparagus bean, Chinese longbean, moth bean, yardlong bean and jackbean) and edible podded beans (including lima bean (green), broad bean (succulent), blackeyed pea, cowpea, and southern pea):

To provide early season protection of seedlings against injury by wireworm, seed corn maggot, bean leaf beetle, imported fire ant, and aphid, apply as a commercial seed treatment at 2 - 4 fl. oz. per hundredweight of seed prior to planting. (Not registered for use in California)

4/4

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

The following plant-back intervals should be observed for listed crops:

ROTATIONAL PLANT-BACK INTERVALS*				
IMMEDIATE PLANT-BACK				30-DAY PLANT-BACK
Barley	Cabbage	Eggplant	Potatoes	Cereals
Beans, edible podded	Canola	Ground cherry	Rape greens	(Including buckwheat, millet, oats, popcorn, rice, rye and triticale)
Beans, succulent shelled	Cauliflower	Kale	Rapeseed	Legumes
Broccoli	Cilantro	Kohlrabi	Sorghum	(Including soybeans, beans [except beans, edible podded and beans, succulent shelled] and peas)
Broccoli, Chinese	Collards	Leaf petiole veg.	Sugar beets	Safflower
Broccoli, raab	Corn, Field	Lettuce	Tomatillo	
Brussels Sprouts	Corn, Sweet	Mustard greens	Tomato	
Cabbage, Chinese mustard	Cotton	Pepinos	Tumip	
Cabbage, Chinese	Cucurbits	Pepper	Wheat	
*Cover crops for soil building or erosion control may be planted any time , but do not graze or harvest for food or feed.				
*All other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient have been established, a 12-month plant-back interval should be observed.				

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with GAUCHO 480 Flowable must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

USER RESTRICTION:

Rape greens grown and harvested from GAUCHO 480 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from GAUCHO 480 treated seed is only for industrial uses and can not be used for edible oil or any other human / feed consumption.

NOTICE - READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS

PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

IN CASE OF AN EMERGENCY ENDANGERING
LIFE OR PROPERTY INVOLVING THIS PRODUCT,
CALL (800) 424-9300

Gustafson (logo) is a Reg. TM of Gustafson LLC
GAUCHO - Reg. TM Bayer AG, Germany
RTU is a Reg. TM of Gustafson LLC
VITAVAX is a Reg. TM of Uniroyal Chemical Co., Inc.
ALLEGIANCE is a TM of Gustafson LLC

A quality product manufactured by:

Gustafson LLC
1400 Preston Road, Suite 400
Plano, Texas 75093

155MLCPSS(S)-105180