

ACCEPTED
 February 23, 2000
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7501-159

7501-159

02/23/2000

1/2

- 1 -

GAUCHO® 75 ST
INSECTICIDE

ACTIVE INGREDIENT:

1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....75.0%
 INERT INGREDIENTS:.....25.0%
 Total.....100.0%

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty container into application equipment by tapping sides to dislodge residues. Then dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.
STATEMENT OF PRACTICAL TREATMENT
 If Swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water. Do not induce vomiting. Do not give anything by mouth to an unconscious person. On Skin: Wash with plenty of soap and water. Get medical attention if irritation develops. If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention. If in Eyes: Flush with plenty of water. Call a physician if irritation develops.
ENVIRONMENTAL HAZARDS
 This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

For Early Season Protection Against Certain Sucking Insects: GAUCHO 75 ST will aid in the protection of seeds and seedlings against injury by early season aphids, chinch bugs and thrips.
Canola:
 To provide early season protection of seedlings against injury by aphids, flea beetle and wireworm apply 10.7 to 21.3 oz. per 100 lbs. of seed prior to planting.
Field Corn:
 For the protection of corn plants from the corn insect pests listed below. Apply as a seed treatment at the recommended rates listed.

PEST	APPLICATION RATE, OZ./CWT. OF SEED
Corn root worms* (including Northern and Western) *(suppression)	10.7
Southern corn root worm Mexican* corn root worm *(suppression)	5.4 - 10.7
Flea beetles Chinch bugs	
Seed corn maggot Wireworms Corn leaf aphid Imported fire ant	2.7 - 5.4
Seed corn maggot Wireworms Imported fire ant	1.4 - 2.7
Wireworms	0.7 - 1.4

EPA Reg. No. 7501-159

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
 For use in commercial seed treaters only. Not for use on agricultural establishments in hopper-box, slurry-box, or other seed treatment applications at, or immediately before, planting.
 This product is to be used in treaters designed for pelleting or coating seed. Mix thoroughly before use or use entire container at one time. All mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

STORAGE AND DISPOSAL
STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.
PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

Wheat, Barley:
 To provide early season protection of seedlings against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly and Wireworms, apply 0.67 to 2.0 oz. per 100 lbs. of seed prior to planting. For wireworm control apply 0.67 to 1.0 oz. per 100 lbs. of seed. Ensure thorough coverage. Use the higher rate to provide increased length of protection and from heavy insect pressure and from wireworms, and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors. Do not graze or feed livestock on treated areas for 45 days after planting.
Sorghum:
 To provide early season protection of seedlings against injury by Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Fire ants and Wireworms, apply 5.3 oz. per 100 lbs. of seed prior to planting. Ensure thorough coverage. Do not graze or feed livestock on treated



areas for 45 days after planting.

Cotton (Delinted Seed Only):

To provide early season protection of seedlings against injury by early season thrips and aphids, apply 4.2 to 5.3 ounces per hundred weight of seed prior to planting. Ensure thorough coverage.

Sugar Beets:

To provide early season protection of seedlings against injury by Whitefly, Aphids, Leafhoppers (including to reduce potential for spread of Curly top and Yellow mosaic virus due to aphid and leafhopper vectors), Root aphid, Thrips and Wireworms, apply as a commercial seed treatment in a water slurry containing 2.0 to 4.25 oz. of GAUCHO 75 ST in or on a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed - approximately 1 kilogram by weight.) Apply in a film coat directly to raw seed (100,000 seed or approximately 1 kilogram by weight.) at the rate of 2.0 oz. per unit. If rates exceed 2.0 oz. per unit, seed must be pelleted. Film coat application and pelleting companies must be approved by Gustafson, Inc.

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

ROTATIONAL PLANT-BACK INTERVALS*				
IMMEDIATE PLANT-BACK				30-DAY PLANT-BACK
Barley	Cabbage	Ground cherry	Potatoes	Cereals (Including buckwheat, corn, millet, oats, popcorn, rice, rye and triticale) Legumes (Including soybeans, beans and peas) Safflower
Broccoli	Canola	Kale	Rape greens	
Broccoli, Chinese	Cauliflower	Kohlrabi	Sorghum	
Broccoli raab	Collards	Lettuce	Sugar beets	
Brussels Sprouts	Cotton	Mustard greens	Tomatillo	
Cabbage, Chinese mustard	Cucurbits	Pepinos	Tomato	
Cabbage, Chinese	Eggplant	Pepper	Wheat	

*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.
*For all other crops not listed on a Imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates. Do not over-treat GAUCHO 480 treated seed with GAUCHO 75 ST.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with GAUCHO 75 ST must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

**NOTICE - READ CAREFULLY
CONDITIONS OF SALE AND LIMITED
WARRANTY STATEMENT**

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC, no claims are made to guarantee germination of carry-over seed. As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in

accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

**IN CASE OF AN EMERGENCY ENDANGERING
LIFE OR PROPERTY INVOLVING THIS PRODUCT,
CALL (800) 424-9300**

A quality product manufactured for:
Gustafson LLC
1400 Preston Road, Suite 400
Plano, Texas 75093
Questions? (972) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson LLC
GAUCHO - Reg. TM Bayer AG, Germany

G75ST(S)-999120