Under the Pederal Insect Pungicide, and Rodenticide Act, as amended, for the pesticide

INSECTICIDE

MPA Rog. No. 75/ ACTIVE INGREDIENT

1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro -2-imidazolidinimine......75.0% INERT INGREDIENTS: 25,0%

KEEP OUT OF REACH OF CHILDREN CAUTION

PECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water. Do not induce vomiting. Do not give anything by mouth to an unconscious person. On Skin: Wash with plenty of soap and water. Get medical attention if irritation develops. If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention. If in Eyes: Flush with plenty of water. Call a physician if irritation develops.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds. Plant end rows last to cover any exposed seeds

EPA Reg. No. 7501-159 EPA Est. No.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only. Not for use on agricultural establishments in hopper-box, slurry-box, or other seed treatment applications at, or immediately before, planting.

This product is to be used in treaters designed for pelleting or coating seed. Mix thoroughly before use or use entire container at one time. All mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

STORAGE AND DISPOSAL

STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse

empty container. Open dumping is prohibited.

~__

Continued next column

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container into application equipment by tapping sides to dislodge residues. Then dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For Early Season Protection Against Certain Sucking Insects: GAUCHO 75 ST will aid in the protection of seeds and seedlings against injury by early season aphids, chinch bugs and thrips. Canola:

To provide early season protection of seedlings against injury by aphids, flea beetle and wireworm apply 10.7 to 21.3 ounces per hundredweight of seed prior to planting.

Field Corn:

For the protection of corn plants from the corn insect pests listed below. Apply as a seed treatment at the recommended rates

PEST	GAUCHO 75 ST		
' = 5 '			
	OZ./ 80,000	02.7 CW1	
	UNIT OF SEED	OF SEED	
Corn root worms*	3.7 -5.0	*******	
(including Northern and			
Western) *(suppression)			
Southern corn root worm		5.4 - 10.7	
Mexican* corn root worm			
*(suppression)			
Flea beetles			
Chinch bugs			
Seed corn maggot		2.7 - 5.4	
Wireworms		2., 0.,	
Corn leaf aphid			
Imported fire ant			
Seed corn maggot		1.4 - 2.7	
Wireworm			
Imported fire ant			
Wireworms		0.7 - 1.4	

Wheat, Barley:

To provide early season protection of seedlings against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly and Wireworms, apply 0.67 to 2.0 ounces per hundredweight of seed prior to planting. Ensure thorough coverage. Use the higher rate to provide increased length of protection and from heavy insect pressure and from wireworms, and to reduce potential spread of Badey yellow dwarf virus due to aphid vectors. Do not graze or feed livestock on treated areas for 45 days after planting.

Sorghum:

To provide early season protection of seedlings against injury by Aphids (including Com leaf, English grain, Greehbug, and Yellow sugar cane aphid), Chinch bugs, Fire ahts and Wireworths, apply 5.3 ounces per hundred weight of seed prior to planting. Ensure thorough coverage. Do not graze or feed livestock on treated areas for 45 days after planting.

Gustafson (





Cotton (Delinted Seed Only):

To provide early season protection of seedlings against injury by early season thrips and aphids, apply 5.3 ounces per hundred weight of seed prior to planting. Ensure thorough coverage.

Sugar Beets:

To provide early season protection of seedlings against injury by Whitefly, Aphids, Leafhoppers (including to reduce potential for spread of Curly top and Yellow mosiac virus due to aphid and leafhopper vectors), Root aphid, Thrips and Wireworms, apply as a commercial seed treatment in a water slurry containing 2.0 to 4.25 ounces of GAUCHO 75 ST in or on a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed - approximately 1 kilogram by weight.) Apply in a film coat directly to raw seed (100,000 seed or approximately 1 kilogram by weight.) at the rate of 2.0 ounces per unit. If rates exceed 2.0 ounces per unit, seed must be pelleted. Film coat application and pelleting companies must be approved by Gustafson, Inc.

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

The following plant-back intervals should be observed for listed crops:

ROTATIONAL PLANT-BACK INTERVALS*						
IMMEDIATE PLANT-BACK			30-DAY PLANT-BACK			
Barley Broccoli Broccoli, Chinese Broccoli raab Brussels Sprouts Cabbage, Chinese mus Cabbage, Chinese	Cabbage Canola Cauliflower Collards Cotton tard Cucurbits Eggplant	Ground cherry Kale Kohlrabi Lettuce Mustard greens Pepinos Pepper	Potatoes Rape greens Sorghum Sugar beets Tomatillo Tomato Wheat	Cereals (Including buckwheat, corn, millet, oats, popcorn, rice, rye and triticale) Legumes (Including soybeans, beans and peas) Safflower		

*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

*For all other crops not listed on a Imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates. Do not overtreat GAUCHO 480 treated seed with GAUCHO 75 ST.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with GAUCHO 75 ST must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

NOTICE - READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC no claims are made to guarantee germination of carry-over seed. As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in

accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL (800) 424-9300

A quality product manufactured for: Gustafson LLC 1400 Preston Road, Suite 400 Plano, Texas 75093 (972) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson LLC GAUCHO - Reg. TM Bayer AG, Germany

G75ST(S)-007130