

7501-161

08/07/2002

ACCEPTED

1/2

162(S)071002 First aid

AUG 7 2002

- 1 -

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7501-161

RAXIL® 0.26FS

Seed Treatment Fungicide

FOR USE ONLY BY COMMERCIAL SEED TREATERS
FOR CONTROL OF SPECIFIED DISEASES ON WHEAT, BARLEY AND OATS

ACTIVE INGREDIENT:

Tebuconazole, Alpha-[2-(4-Chlorophenyl)ethyl]-alpha-(1,1-dimethyl-ethyl)-1H-1,2,4-triazole-1-ethanol 3%

INERT INGREDIENTS: 97%

TOTAL 100%

Contains 0.26 lbs. tebuconazole per gallon

STOP - READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 7501-161 EPA Est.
NET CONTENTS:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Avoid breathing vapor or spray mist. Wear long sleeved shirt, long pants or coveralls, and chemical-resistant gloves made of waterproof material. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

FIRST AID

Have product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed:

-Call a poison control center or doctor immediately for treatment advice.

-Have person sip a glass of water if able to swallow.

-Do not induce vomiting unless told to do so by poison control center or doctor.

-Do not give anything by mouth to an unconscious person.

If on skin or clothing:

-Take off contaminated clothing.

-Rinse skin immediately with plenty of water for 15 - 20 minutes.

-Call a poison control center or doctor for treatment advice.

Continued Next Column

If in eyes:

-Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.

-Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

-Call a poison control center or doctor for treatment advice.

If inhaled:

-Move person to fresh air.

-If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

-Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

IMPORTANT: Read these entire Directions and Warranty before using RAXIL 0.26FS.

RAXIL controls or suppresses certain designated seed and soilborne diseases of wheat, barley and oats.

SEED TREATMENT: It is recommended that RAXIL 0.26FS be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9 to 20 fl. oz. of water per 100 lb of seed. Add dye** to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

**21CFR Chapter I, Section 2.25 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored with an EPA approved dye, such as one of the dyes listed in 40 CFR Section 180.1001 (c) and (d), to prevent their subsequent inadvertent use as a food for man or feed for animals.

Gustafson



APPLICATION RATE		
CROP & DISEASE	RATE OF RAXIL 2.6F	APPLICATION DIRECTIONS
<p>Wheat: Stinking Smut, Flag Smut, Loose Smut, Early Season <i>Septoria</i> Disease Complex, Early Season, <i>Rhizoctonia</i> Root Rot, Early Season Common Root Rot, Early Season <i>Fusarium</i> Foot Rot, Early Season, Suppression of Powdery Mildew, Early Season, Suppression of Wheat Leaf Rust</p> <p>Barley: Covered Smut, Loose Smut, Barley Stripe (suppression), Early Season <i>Rhizoctonia</i> Root Rot, Early Season Common Root Rot, Early Season, <i>Fusarium</i> Foot Rot, Early Season Suppression of Powdery Mildew Early Season Suppression of Barley Leaf Rust</p> <p>Oats: Loose Smut, <i>Pythium</i> Damping-off, Early Season <i>Rhizoctonia</i> Root Rot, Early Season Common Root Rot, Early Season <i>Fusarium</i> Foot Rot, Early Season Suppression of Powdery Mildew, Early Season Suppression of Crown rust</p>	1 fluid ounce per 100 pounds of seed	Refer to the SEED TREATMENT section on this label for the recommended method of treating seed.

SEED LABELING REQUIREMENTS

To meet U.S. Federal Seed Act requirements, all seed treated with RAXIL 0.26FS should be labeled: "Treated with alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol (RAXIL). Do not use treated seed for feed, food or oil purposes. Barley, wheat and oat green forage may be grazed or harvested for hay 31 days after seeding."

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Contact Gustafson or your chemical supplier for container return instructions. If container is not returned in accordance with instructions, triple rinse (or equivalent). Then offer for recycling or reconditioning, or, puncture and dispose in a sanitary landfill, or incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE-READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical

treatment. Due to seed quality conditions beyond the control of Gustafson LLC, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC, SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product manufactured by:
Gustafson LLC

1400 Preston Road, Suite 400
Plano, Texas 75093

Questions? (972) 985-8877

RAXIL is a registered trademark of BAYER AG, Germany
Gustafson (logo) is a registered trademark of Gustafson LLC Plano, Texas