

PM 0.1

7501-168

8/26/97 08/97

1/3

RAXIL® 2.6 F

RECD EPA/OPP/DPD1

Seed Treatment Fungicide

'97 MAY 14 AIO:10

FOR USE ONLY BY COMMERCIAL SEED TREATERS

FOR CONTROL OF SPECIFIED DISEASES ON WHEAT, BARLEY, AND OATS

ACTIVE INGREDIENTS:

Tebuconazole, Alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol	28.3%
INERT INGREDIENTS	71.7%
TOTAL	100.0%

Contains 2.6 lb tebuconazole per gallon.

STOP - READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 7501-168

EPA Est No.

Net Contents:

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF EYES ARE CONTAMINATED: Hold eyelids open and flush with plenty of water for 15 minutes. Call a physician if irritation persists or develops after flushing.

IF INHALED: First remove victim to fresh air or uncontaminated area. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention as soon as possible.

SYMPTOMS OF POISONING: The compound does not cause

(Cont. Next Column)

any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

IMPORTANT: Read these entire Directions and Warranty before using RAXIL 2.6 F.

RAXIL controls or suppresses certain designated seed and soilborne diseases of wheat, barley and oats.

SEED TREATMENT: It is recommended that RAXIL 2.6 F be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9 to 20 fl. oz. of water per 100 lb of seed. Add dye** to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

**21 CFR Chapter I, Section 2.25 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored with an EPA approved dye, such as one of the dyes listed in 40 CFR Section 180.1001 (c) and (d), to prevent their subsequent inadvertent use as a food for man or feed for animals.

NOTE TO USER:

For worker protection during mixing, wear a long sleeve shirt, long legged trousers or overalls, and chemical-resistant gloves.

Gustafson (logo) is a Reg. TM of Gustafson, Inc.

RAXIL is a Reg. TM used under license.

A quality product manufactured by:

GUSTAFSON, INC.

1400 Preston Road

Suite 400

Plano, Texas 75093

(972) 985-8877

R2.6F(S) 794240



RECOMMENDED APPLICATIONS			
CROP	DISEASE	RATE OF RAXIL 2.6 F	REMARKS
Wheat Barley Oats	<p><u>Wheat:</u> Stinking Smut Flag Smut Loose Smut Early Season Septoria Disease Complex Early Season Rhizoctonia Root Rot Early Season Common Root Rot Early Season Fusarium Foot Rot Early Season Suppression of Powdery Mildew Early Season Suppression of Wheat Leaf Rust</p> <p><u>Barley:</u> Covered Smut Loose Smut Barley Stripe (suppression) Early Season Rhizoctonia Root Rot Early Season Common Root Rot Early Season Fusarium Foot Rot Early Season Suppression of Powdery Mildew Early Season Suppression of Barley Leaf Rust</p> <p><u>Oats:</u> Loose Smut Pythium Damping-Off Early Season Rhizoctonia Root Rot Early Season Common Root Rot Early Season Fusarium Foot Rot Early Season Suppression of Powdery Mildew Early Season Suppression of Crown Rust</p>	0.1 fluid ounces per 100 pounds of seed	Refer to the SEED TREATMENT section on this label for the recommended method of treating seed.

SEED LABELING REQUIREMENTS

To meet U. S. Federal Seed Act requirements, all seed treated with RAXIL 2.6 F should be labeled: "Treated with alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol (RAXIL). Do not use treated seed for feed, food or oil purposes." Do not graze barley, wheat, or oat green forage before 31, 38, and 51 days after seeding respectively.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE - READ CAREFULLY
CONDITIONS OF SALE AND LIMITED WARRANTY
STATEMENT**

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson, Inc., warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON, INC. NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson, Inc. If these terms are not acceptable, return all products to the place of purchase; unopened for a full refund.

