1/4

	on reverse before comple	ntina form	,			D80, Approvel expires 2-28-9	
≎EPA	Environmenta	United States		X	Registration Amendment Other	OPP Identifier Number	
		Application	n for Pestici	de - Section	1		
1. Company/Product Nu	mber		2. EPA	Product Manager	, 3.	Proposed Classification	
750/	-/7/ ame)		She	eradu Ho.	bgood [None Restricted	
RAXIL XT W	ettable Powder	Fungica	de 21	No tificati	ons		
5. Name and Address of Applicant Include ZIP Code) Gustafson LLC 1400 Preston Rd., Suite 400 Plano, TX 75093				6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.			
Check it	this is a new address		Produ	ct Name			
			Section - I				
			Oection - I				
Amendment - Explain below.				Final printed labels in repsonse to NOTIFICATION Agency letter dated			
Resubmission in response to Agency letter dated							
Notification - Explain below.				— Me Too" Application. APR 2 3 2002 Other - Explain below.			
the 1/14/06 California Seed treating	uce required EPA stam - Only for use gestablisher use,	red lab in communents, No	el, "IN nercial otallowed Section - II	for_	her understand that if this notificati to 98-16 and 40 CFR 152-46, this pe be subject to unforcement action as	ridy make any false statement to EPA. I look to not considered with the forms of PR volution by the property of the volution of PTPRA, and I id possible under sections 12 and 14 of the fall of the f	
1. Material This Product	Will Be Peckaged In:				.,		
Child-Resistant Packagin	ng Unit Packaging		Water Soluble P	sckaging	2. Type of Contain		
Yes	Yes		Yes		Metal Plastic		
No	No No		No No		Glass	•	
* Certification mus be submitted	f "Yes" Unit Packaging wgt	No. per . container	if "Yes" Package wgt	No. per container	Paper Other	(Specify)	
3. Lacation of Net Conte	Container	4. Size(s) Reta	il Container	5. La	ocation of Label Direc	tions	
6. Manner in Which Label is Affixed to Product		Lithograph OtherStenciled					
			Section - I\	/			
1. Contact Point (Comp	lete items directly below	for identification	of individual to be	contacted, if nec	essary, to process th	is application.)	
Name Dennis McFadden Mgs				& Keg Afy		one No. (Include Area Code)	
	tatements I have made or it any knowlingliy false or able law.	Certificati n this form and a	ion Il attachments the	reto are true, acc	urate and complete.	6. Date Application Received (Stamper)	
2. Signature		3	, Tide				
Denne A	1 chadden		Algr. Tox	BReg Af	fairs		
4. Typed Name	2	5	. Date	/ /			
Drunic L	1-6-220		4	117/02			

NOTIFICATION

APR 23 2012

RAXIL®XT

Wettable Powder Fungicide

ACTIVE INGREDIENT:

Tebuconazole [Alpha-[2-(4-Chlorophenyl)ethyl]-	
alpha-(1,1-dimethyl-ethyl)-1H-1,2,4-triazole-1-	
ethanol]	15.0%
Metalaxyl [N-(2,6-dimethylphenyl)-N-	
(methoxyacetyl)alanine methyl ester]	20.0%
INERTINGREDIENTS:	
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 7501-171 EPA Est. Net Contents:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND **DOMESTIC ANIMALS** CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling. Causes eye injury (irritation). Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear; long-sleeved shirt and long pants and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- -Call a poison control center or doctor immediately for treatment advice.
- -Have person sip a glass of water if able to swallow.
- -Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything by mouth to an unconscious person.

- -Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treament advice.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish, estuarine and marine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. PESTICIDE DISPOSAL: This pesticide's wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control agency, or the Hazardous Waste Representative at the nearest EPA Re | gional Office for guidance.

Gustafson





CONTAINER DISPOSAL: When all water soluble bags are used, the outer package should be clean and may be disposed of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. If the outer container contacts formulated product in any way, it must be triple rinsed with clean water. Add rinsate to the mix tank and dispose of the outer package as described above.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the areas during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultrual pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material and shoes plus socks.

RAXIL XT Wettable Powder Fungicide is a combination of systemic fungicides. Tebuconazole and Metalaxyl aid in the control or suppression of the following seed, seedling and soilborne diseases of wheat, barley and oats:

Wheat - Stinking smut, Flag smut, Loose smut, Early season Septoria disease complex, Early season Rhizoctonia root rot, Early season common root rot, Early season Fusarium foot rot, Early season suppression of powdery mildew, Early season suppression of wheat leaf rust, Seedborne Fusarium scab, General seed rots and Pythium damping-off;

Barley - Covered smut, Loose smut, Barley stripe (suppression), Early season Rhizoctonia root rot, Early season common root rot, Early season Fusarium foot rot, Early season suppression of powdery mildew, Early season suppression of barley leaf rust, Seedborne Fusarium scab, General seed rots and Pythium damping-off;

Oats - Loose smut, Early season Rhizoctonia root rot, Early season common root rot, Early season Fusarium foot rot, Early season suppression of powdery mildew, Early season suppression of crown rust, Seedborne Fusarium scab, General seed rots and Pythium damping-off.

Consult your area agronomist regarding planting depth recommendations for the particular variety to be planted.

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

RECOMMENDATIONS FOR USE

The pouches of RAXIL XT Wettable Powder are water soluble. Do not allow pouches to become wet prior to adding to the mix tank. Do not handle pouches with wet hands or wet gloves. Do not cut or puncture the pouches. Use the entire water-soluble bag. Do not break pouches or use partial pouches.

Prepare no more slurry than is needed for immediate treating. Mechanical agitation is required for proper mixing of RAXIL XT Wettable Powder. Agitate thoroughly before and during application. Add all of the required water to the mix tank. With agitation running, drop the required number of pouches into the tank. If this product is packaged with separate colorant pouches, add one colorant pouch with each RAXIL XT Wettable Powder pouch used. One 0.5 pound pouch of RAXIL XT Wettable Powder and one 0.5 pound pouch of dry colorant should be used to treat 40 50 hundredweights of seed. This equates to a use rate of from 0.16 - 0.20 ounces per hundredweight of seed and will deliver from 15 - 18.75 parts per million of tebuconazole and from 20 -25 parts per million of metalaxyl. (One full case of product can treat from 960 - 1200 hundredweights of seed). Product should be diluted with sufficient water to give an applied slurry rate of 9 to 20 fluid ounces per 100 pounds of seed. Applications should be made using standard slurry or mist-type seed treatment equipment. Uniform application to seed is necessary to ensure seed safety and best disease control. Seed should be sound and well cured prior to treatment.

NOTE: This product is undyed. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR. Part 2.25. Any dye added to treated seed must be cleared for use

To meet U.S. Federal Seed Act requirements, all seed treated with RAXIL XT Wettable Powder should be labeled:

TREATED SEED.

DONOTUSE FOR FOOD, FEED OR OIL PURPOSES.

Treated with Alpha-[2-(4-Chlorophenyl)ethyl]-alpha-(1,1dimethylethyl)-1H-1,2,4-triazole-1-ethanol (Tebuconazole) and Metalaxyl.

Seed labeling must contain the following restriction: "Barley, wheat and oat green forage may be grazed or harvested for hay 31 days after seeding."

NOTICE-READCAREFULLY **CONDITIONS OF SALE AND LIMITED WARRANTY** STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC no claims are made to guarantee germination of carry-over

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES SUBJECT TO APPLICABLE LAW, GUSTAFSON LICNEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRE-SENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use and store this product in accordance with the safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing

Conditions of Sale and Limited Warranty, which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

A quality product packaged for:

Gustafson LLC 1400 Preston Road, Suite 400 Plano, Texas 75093 Questions? (972) 985-8877 RAXIL is a Reg. TM of Bayer, A.G., used under license Gustafson (logo) is a Reg. TM of Gustafson LLC