c0.87.	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs	EPA Reg. Number:	Date of Issuance:	
	Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460	7501-197	JAN 31200	
NOTICE OF PESTICIDE: <u>x</u> Registration under FIFRA, as amended)		Term of Issuance:	Term of Issuance: Conditional	
		Name of Pesticide Product: Imidacloprid Metalaxyl Seed Treatment		
Name and Address of Registrant (include Gustafson LLC 400 Preston Rd., Suite 4 Plano, TX 75093 attn: Dennis McFadden				
On the basis of information furnished by Rodenticide Act. Registration is in no way to be construed Administrator, on his motion, may at any	any correspondence on this product always refer to the above f the registrant, the above named pesticide is hereby registered/re as an endorsement or recommendation of this product by the A time suspend or cancel the registration of a pesticide in accorda his Act is not to be construed as giving the registrant a right to es	registered under the Federal l gency. In order to protect he nuce with the Act. The accep	alth and the environment, the tance of any name in connecting	
pesticide is registered, he eliminate the need for co time, additional data are require submission of suc	tionally registered in accordance with owever, it is not regarded as permaner ontinual reassessment of pesticides. If required to maintain in effect an existi ch data under FIFRA section $(3)(c)(2)$ egistration Number to read, "EPA Reg	the Agency deter ng registration, the (B).	Registration does n mines that, at any ne Agency will	
Signature of Approving Official:		Date:		
11.		JAN 31	2003	
/ Dani Daniel Insecticide-Rodenticide Branch	1			

ļ

1/4

Page Two

.....

7501-197

2. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

、

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact me at (703) 305-5409.

Enclosure :

IMIDACLOPRID-METALAXYL Seed Treatment

For the protection of Seeds and Seedlings of Field Corn, Sweet Corn and Popcorn against Damping-Off, Seed Decay, Seedling Blight; also against Seed Corn Beetle, Seed Corn Maggots, and Wireworms.

ACTIVE INGREDIENTS:

Imidacloprid (1-[(6-chloro-3-pyridinyl)methyl]-N-nitro-2-Imidazolidinimine)	
Metalaxyl (N-2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester 1.00%	
INERT INGREDIENTS	
TOTAL	

EPA Reg. No. 7501-EPA Est.

NET CONTENTS:

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear: longsleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED with COMMENTS In EPA Letter Dated:

JAN 3 | 2003

Under the Fungicide	Federal and Rod	l Insect lenticid	icido, o Act,
es amend	ed. for	the pes	CHO CHO CHO
registered		2	2

Manufactured for:



Gustafson LLC 1400 Preston Rd., Suite 400 Plano, TX 75093 Questions? (972) 985-8877 **USER SAFETY RECOMMENDATIONS:** Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID	
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person.
if in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
	 Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL:

CHEMTREC: (800) 424-9300 (Toll free – US) CHEMTREC: (703) 527-3387 (Domestic and International)

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates and toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Exposed treated seeds may be hazardous to birds. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. Do not contaminate water when disposing of equipment washwaters or rinsate.

PRODUCT INFORMATION

Imidacloprid-Metalaxyl may be used on seed previously treated with a full dosage of protective fungicide to give added protection against *Pythium* damping-off or seed decay. **Note:** Adjust the seed drill or planter to give the required rate of seed delivery with treated seed. Regularly inspect planter plates and clean as required. At the end of each day, clean the electric-eye monitors as directed by the manufacturer. **Important:** Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seed beyond planting time.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

4/4

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, protective eyewear.

Field Corn, Sweet Corn, Popcorn: Use at a rate of 1.5 ounces per 42 pounds of seed.

Mixing Instructions: (1.5 oz. canister) Use as a dry mixture in the planter box as a seed treatment prior to planting. Apply Imidacloprid-Metalaxyl to seed at planting time with the canister applicator tube system, using one 1.5 oz. canister per 42 pounds of seed. For best results, fill planter box with seed, add Imidacloprid-Metalaxyl through applicator tube and mix so that all seeds are covered. DO NOT MIX WITH BARE HANDS. Use only at the recommended rate. Lower amounts may not give desired control. Excessive amounts may cause seed injury.

Mixing instructions: (4-pound pail) Field corn, Sweet Corn, Popcorn: Use two scoops full (1.5 oz.) per 42 pounds of seed.

Use as a dry mixture in the planter box as a seed treatment prior to planting. For best results, fill planter box half full of seed, add half the required amount of Imidacloprid-Metalaxyl and mix thoroughly with a stick or paddle. Add remainder of the seed and Imidacloprid-Metalaxyl and mix again so that all seed are covered. DO NOT MIX_WITH BARE HANDS. Use only at recommended rate. Lower amounts may not give desired control Excessive amounts may cause injury.

ROTATIONAL CROPS¹, Treated areas may be replanted with any crop specified on the Imidacloprid-Metalaxyl label, or for any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. The following plant-back intervals should be observed for listed crops:

Popcorn	Cereals
Potatoes Rape greens Rapeseed Sorghum Sugar beets Tomatillo Tomato Tumip greens Wheat	(Including buckwheat, millet, oats, popcom, rice, rye and triticale) Safflower Soybeans
	Rapeseed Sorghum Sugar beets Tomatillo Tomato Turnip greens

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry location. Protect from excessive heat.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty canister into application equipment. Then dispose of empty canister in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE - READ CAREFULLY

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases[§]this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

Gustafson (logo) is a Reg. TM of Gustafson LLC, Plano, Texas GAUCHO[®] is a registered trademark of Bayer AG, Germany