

NET WEIGHT 5 LBS. HOBBY'S

1537-2

ACCEPTED
Jul 19 1967
U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
WASHINGTON, D.C. 20250

PROLIN* READY - TO - USE RAT & MOUSE BAIT NEW IMPROVED WARFARIN RODENTICIDE

**For hard-to-kill rats and mice. A better and more effective rodenticide.
For increased effectiveness. The most notable advance in rodenticides since
Warfarin.**

The antibacterial agent increases the effectiveness of the product and assures more thorough control of rats and mice. Especially recommended for rat and mouse infestations which are difficult to control.

ACTIVE INGREDIENTS

N-(2-Quinoxaliny) sulfanilamide	0.025%
Warfarin 3-acetonylbenzyl-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.950%
TOTAL	100.00%

*Prolin is a Trademark of the Wisconsin Alumni Research Foundation
Prolin Warfarin plus an antibacterial agent is manufactured under license from the Wisconsin Alumni Research Foundation, U.S. Patent No. 2,427,578, other patent applications pending.

DIRECTIONS FOR USE:

RATS Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

CAUTION KEEP OUT OF REACH OF CHILDREN

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hem-

N-(2-Quinoxalinyloxy) succinimide	0.025%
Warfarin 3-(acetylbenzyl)-4-hydroxycoumarin ..	0.025%
INERT INGREDIENTS	99.950%
TOTAL	100.00%

*Prolin is a Trademark of the Wisconsin Alumni Research Foundation.
 Prolin Warfarin plus an antibacterial agent is manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,411,075. See patent applications pending.

DIRECTIONS FOR USE:

RATS—Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

MICE—Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

**CAUTION
 KEEP OUT OF REACH OF
 CHILDREN**

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin. It hereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

USDA—REG. NO. 7537-2

NET CONTENTS 5 LBS.

Manufactured By:

MONROE GRAIN & SUPPLY, INC.

MONROE, INDIANA