

7537-2

08/12/2004

received 6/14/2004

1/1

# Hobby's

READY TO USE  
**RAT AND MOUSE BAIT**  
KILLS NORWAY RATS, ROOF RATS, AND HOUSE MICE



**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 2.5 lb. (1.134 Kg.)

## Hobby's



**ACCEPTED**  
AUG 12 2004  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7537-2  
EPA ESTABLISHMENT NO. 7537-01-1  
EPA REGISTRATION NO. 7537-2

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL**-Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides.

To help prevent accidents:

1. Store product not in use in a location out of reach of children and other pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are filled, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

**USE RESTRICTIONS-**For control of Norway rats, roof rats, and house mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.  
**SELECTION OF TREATMENT AREAS-**Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS

**RATS:** Apply 4 oz. of bait at single location. This package only contains enough bait for 1 to 2 rats. Additional packages of bait may be needed. -Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8-12 foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish bait stations and replenish as needed.

**ACTIVE INGREDIENT-**  
WARFARIN 0.025%  
**INERT INGREDIENTS** 99.975%  
**TOTAL** 100.00%

**PRECAUTIONARY STATEMENTS**  
**-HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
Caution: Keep away from humans, domestic animals and pets. If bait is handled, wear waterproof gloves. After handling bait, wash hands thoroughly.  
**ENVIRONMENTAL HAZARDS**  
This product is toxic to mammals and birds. Do not apply directly to water.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed	•Call poison control center or doctor immediately for treatment advice or transport the patient to the nearest hospital. •Do not drink water. •Do not administer anything by mouth or induce vomiting unless told to do so by the poison control center or doctor.
If on skin or clothing	•Wash hands thoroughly with soap and water.
NOTE TO PHYSICIAN	
If swallowed, Warfarin may reduce the clotting time of the blood and cause bleeding. If ingested, administer Vitamin K intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	

**STORAGE AND DISPOSAL**  
**Store:** Store only in original container, in a cool, dry place inaccessible to children and pets.  
**Disposal:** Do not reuse this container. Place in trash or offer for recycling as available. If partly filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.  
Manufactured by: **HOWBERT CO.**  
P.O. BOX 88, MT. HOPE, OHIO 44660  
800-213-7562