

Germicide . . . Udder Wash . . . Milkstone Solvent . . . Drying Agent

U. S. I O D O P H O R

FORMULA NO. 34

SANITIZER — DETERGENT

USDA REG No. 7546-1

MANUFACTURED BY

United States Chemical Corporation

MILWAUKEE, WISCONSIN

Net Contents: U. S. Gallon

Active Ingredients:

Nonylphenoxypoly (ethyleneoxy) ethanol-iodine complex (providing 1.75% available iodine).....	13.0%
Phosphoric Acid	8.0%
Inert Ingredients	79.0%
Total	100.0%

ACCEPTED
6-8-65
UNDER THE FEDERAL INSECT PESTICIDE ACT
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. 7546-1

CAUTION: HARMFUL IF SWALLOWED.
KEEP OUT OF CHILDREN'S REACH.
SEE LEFT PANEL — READ PRECAUTIONS CAREFULLY.

"Sanitation . . . For The Nation!"




UDDER WASHING

1. Use fresh towel or sponge for each cow.
2. Wash udder and teats with warm 25 ppm solution of **U. S. IODOPHOR**. Discard towel.
3. Check each quarter with strip-cup.
4. Use teat-cup with 25 ppm **U. S. IODOPHOR** solution to protect each teat-canal and discourage mastitis infections.

UTENSILS AND EQUIPMENT

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a detergent solution of **U. S. TALENT** or **ORBIT**.
3. Rinse with clean water.
4. Sanitize in a **U. S. IODOPHOR** solution of 1 oz. per 5 gallons of water (25 ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a **CLEAN, SANITIZED** rack or drainboard to air dry.

For best results, use tepid or lukewarm solutions (room temperature up to bath-water temperature) 70° to 110° F. Use at recommended strengths.



PROPORTIONS TO USE

For 50 ppm, use 1 oz. **U. S. IODOPHOR** to 2½ gallons water.

For 25 ppm, use 1 oz. **U. S. IODOPHOR** to 5 gallons water.

CAUTION: **U. S. IODOPHOR** is a concentrated material. Causes skin irritation. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. Get medical attention for eyes.

Avoid contamination of foods.

3. Check each quarter with strip-cup.
4. Use teat-cup with 25 ppm U. S. IODOPHOR solution to protect each teat-canal and discourage mastitis infections.

UTENSILS AND EQUIPMENT

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a detergent solution of U. S. TALENT or ORBIT.
3. Rinse with clean water.
4. Sanitize in a U. S. IODOPHOR solution of 1 oz. per 5 gallons of water (25 ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a **CLEAN, SANITIZED** rack or drainboard to air dry.

For best results, use tepid or lukewarm solutions (room temperature up to bath-water temperature) 70° to 110° F. Use at recommended strengths.

ACCEPTED

6-8-65

UNDER THE FEDERAL INSECTICIDE
REGISTRATION ACT
FOR ECONOMIC CONTROL



PROPORTIONS TO USE

For 50 ppm, use 1 oz. U. S. IODOPHOR to 2½ gallons water.

For 25 ppm, use 1 oz. U. S. IODOPHOR to 5 gallons water.

CAUTION: U. S. IODOPHOR is a concentrated material. Causes skin irritation. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. Get medical attention for eyes.

Avoid contamination of foods.

Use at recommended strengths and at temperatures not above 110° F.



ACCEPTED
 6-8-65
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 7546-1

IODOPHOR

FORMULA NO. 34

SANITIZER — DETERGENT

USDA REG No. 7546-1

MANUFACTURED FOR
PURE MILK ASSOCIATION
 CHICAGO, ILLINOIS

Net Contents: 1 U. S. Gallon

Active Ingredients:

Nonylphenoxypoly (ethyleneoxy) ethanol-iodine complex (providing 1.75% available iodine)	13.0%
Phosphoric Acid	8.0%
Inert Ingredients	79.0%
Total	100.0%


CAUTION: AVOID CONTAMINATION OF FOODS.
 KEEP OUT OF CHILDREN'S REACH.
 HARMFUL IF SWALLOWED.
 SEE SIDE PANEL — READ PRECAUTIONS
 CAREFULLY.

UTENSILS AND EQUIPMENT

1. Prewash utensils.
2. Wash with a **PMA** detergent solution.
3. Rinse with clean water.
4. Sanitize in **PMA IODOPHOR No. 34** solution of 1 oz per 5 gallons of water (25 ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a **CLEAN, SANITIZED** rack to air dry.


For best results, use tepid or lukewarm solutions (room temperature up to bath-water temperature) 70° to 110° F. Use at recommended strengths. Solution must be discarded when concentration falls below 12½ ppm.

CAUTION: PMA IODOPHOR No. 34 is a concentrated material. Causes skin irritation. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. Get medical attention for eyes.



UTENSIL WASHING

Causes skin irritation. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. Get medical attention for eyes.



UDDER WASHING

1. Use fresh towel or sponge for each cow.
2. Wash udder and teats with warm 25 ppm solution of **PMA IODOPHOR No. 34**. Discard towel.
3. Check each quarter with strip-cup.

PROPORTIONS TO USE

For 50 ppm, use 1 oz. **PMA IODOPHOR No. 34** to 2½ gallons water.

For 25 ppm, use 1 oz. **PMA IODOPHOR No. 34** to 5 gallons water.

Solution must be discarded when concentration falls below 12½ ppm.

lution

For 200 ppm* titratable chlorine,
Use 1 oz. per 3 gals. water.

of available chlorine

CAUTION:

*ppm means "parts per million".



KING OF MODERN SAN