


MODERN SANITIZERS

7546-2

on

N

NGS



DAIRY USE INSTRUCTIONS

Preclean all milking equipment and utensils.

- 1. Wash with a strong detergent solution.
- 2. Rinse with clean water.
- 3. Sanitize with a 100 ppm SAN-I-KING solution.
- 4. Rinse with clean water.
- 5. Sanitize with a 200 ppm SAN-I-KING solution just before next use.

DAIRY AND FOOD PLANTS

Sanitize all precleaned milk and food contact equipment just before use with a 200 ppm SAN-I-KING solution.

CAUTION: Do not use with acid detergents.

Cameraman's Note

Poor Copy

REGISTRATION



INSTITUTIONAL USE DIRECTIONS

UTENSIL SANITIZATION

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash in a U.S. detergent solution.
3. Rinse with clean water.
4. Sanitize in a SAN-I-QUEEN solution of 1 oz. to 2 gallons of water (200 p.p.m.). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a CLEAN, SANITIZED rack or drainboard to air dry.

TABLE OF PROPORTIONS

For 200 ppm available chlorine, use 1 oz. per 2 gallons water.

1 ppm means 1 part per million.

CAUTION: Do not mix SAN-I-QUEEN with acid detergents.

Directions: For sanitizing, use 1 oz. per 2 gallons of water.

ANTIDOTE:

For eye contact, flush with plenty of water for at least 15 minutes.

For skin contact, flush with plenty of water for at least 15 minutes.


CALL A PHYSICIAN

7239

Cameraman's Note

Poor Copy

754



SAN-I-QUEEN

LIQUID SODIUM HYPOCHLORITE

E.P.A. Reg.

No. 80

"QUEEN OF MODERN SANITIZERS"

MANUFACTURED BY

United States Chemical Corpora

MILWAUKEE, WISCONSIN 53215

NET CONTENTS: 1 U.S. GALLON

Active ingredient - Sodium hypochlorite
Inert ingredients

CAUTION: KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
IRRITATING TO EYES, SKIN AND LUNGS
SEE SIDE PANEL FOR ANTIDOTE

"Sanitation . . . For The Nation"