USE DIRECTIONS

DISINFECTION

Use at 1 oz. per gallon of water. This product will be effective against Pseudomonas aeruginosa PRD-10 at this level.

At 1 oz. per gallon of water, this product is fungicidal against pathogenic fungi.

For best results, use EMERGE as directed above with a cloth or mop and apply to walls, floors, and other hard surfaces.

For heavy soil or organic matter, a pre-cleaning step is recommended.

SANITIZATION

If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm* (1 oz. per 3.5 gallons of water). NO RINSE REQUIRED AT THIS LEVEL.

This product when used on environmental, inanimate, hard surfaces at 1 oz. per gallon of water, is effective against Influenza A_2, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses.

*ppm means "parts per million" of quaternary ammonium chlorides.

Phenol Coefficients
Guaranteed Minimum
S. aureus   78
S. typhosa   48

NET CONTENTS: U. S. GALLON

---

EMERGE™

No. 75

EPA Reg. No. 7546-18
EPA EST. NO. 7546-WI-1

CONCENTRATED DETERGENT, SANITIZER,
FUNG` `, DISINFEC` `, DEODORIZER

For

FOOD SERVICE MAINTENANCE CLEANING

Spray, Brush, or Soak

ACTIVE INGREDIENTS:

n-Alkyl (60% C_{14}, 30% C_{16}, 5% C_{12}, 5% C_8)
  dimethyl benzyl ammonium chlorides ..   4.5%
n-Alkyl (68% C_{12}, 32% C_{14}) dimethyl
  ethylbenzyl ammonium chlorides 4.5%
Tetrasodium ethylenediamine tetraacetate   2.0%
Sodium Carbonate 4.0%
INERT INGREDIENTS: 85.0%

                                 100.0%

DANGER

Keep Out of Reach of Children. See right side panel for additional precautionary and first aid statements.

ACCEPTED

JUL 6 1976

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
EPA Reg. No.
registered under 7546-18

EMERGE™

No. 75

EPA Reg. No. 7546-18

EPA EST. NO. 7546-WI-1

CONCENTRATED DETERGENT, SANITIZER,
FUNG⋅ E, DISINFEC⋅ T, DEODORIZER

For

FOOD SERVICE MAINTENANCE CLEANING

Spray, Brush, or Soak

ACTIVE INGREDIENTS:

n-Alkyl (60% C ⋅, 30% C ⋅, 5% C ₍₂, 5% C ₍₈)
   dimethyl benzyl ammonium chlorides . . 4.5%
n-Alkyl (68% C ₍₂, 32% C ₍₄) dimethyl
   ethylbenzyl ammonium chlorides 4.5%
Tetrasodium ethylenediamine tetraacetate 2.0%
Sodium Carbonate 4.0%
INERT INGREDIENTS: 85.0%

                                 100.0%

DANGER

Keep Out of Reach of Children. See right side panel for additional precautionary and first aid statements.

MANUFACTURED BY

United States Chemical Corporation

P.O. BOX 366   WATERTOWN, WIS. 53094

GENERAL INFORMATION

EMERGE is a concentrated detergent, disinfectant, sanitizer, fungicide, and deodorizer. It is specifically designed for Institutional use, such as Hospitals, Nursing Homes, and Schools where housekeeping will be of prime importance in controlling the hazard of cross-infection from environmental surfaces.

EMERGE is for use on floors, walls, and other hard surfaces in Hospitals, Nursing Homes, Hotels, Public Areas, Industrial Areas, and in Institutions. It is excellent for Operating Rooms, Rest Rooms, Nurseries, and Food Services Areas.

When used as directed, EMERGE is effective against Staphylococcus a u r e u s. Salmonella choleraesuis, and Pseudomonas aeruginosa. EMERGE is also effective against pathogenic fungi, Trichophyton interdigitale.

If used for sanitization of previously cleaned food processing equipment or food contact items, limit the active quaternaries to 200 ppm. At this level, NO POTABLE WATER RINSE is required.

DANGER

Keep Out of Reach of Children. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.