

SANCO-PHENE

March 5, 1973
EPA REG. NO. 7594-6

Comment

USE DIRECTIONS:

Use at 1:100 dilution (approx. 1¼ oz. per gallon of water). Sanco-Phene will provide 850 ppm phenols to this suggested use solution. This will provide disinfection against pathogenic organisms listed below, and consequently qualifies as a disinfectant for hospital use. For heavily soiled areas, use 2½ oz. per gallon of water.

Staphylococcus aureus

Sixty replicates on three preparations — also thirty replicates each on duplicate samples after sixty-day shelf-life A.O.A.C. Method.

Salmonella Choleraesuis

Thirty replicates each on three preparations

Pseudomonas Aeruginosa

Thirty replicates each on three preparations

Manufactured By •

**Sanitary** PRODUCTS CORP.
88 FARWELL STREET • WEST HAVEN, CONN. 06516
U.S.A.

CLEANER DISINFECTANT

DEODORANT

ACTIVE INGREDIENTS

o-Benzyl-p-chlorophenol	4.00%
o-Phenylphenol	3.00%
p-Tertiary amyphenol	1.50%
Tetrasodium ethylenediamine-tetraacetate	1.11%
Sodium Dodecylbenzenesulfonate	6.30%
Sodium xylene sulfonate	3.60%
Isopropanol	19.80%
INERT INGREDIENTS*	50.89%
	100.00%

*Contains cleaning and water softening agents. Sanco-Phene is a concentrated, phosphate free, germicidal cleaner intended for cleaning and disinfection. It is designed for cleaning floors, walls, bed frames, furniture, and bathroom fixtures in a wiping, mopping, or scrubbing operation to wet surfaces thoroughly with the recommended use dilution.

It is free-rinsing and does not leave a film. Consequently, it is recommended for the cleaning of conductive flooring in operating rooms.

NET CONTENTS _____

EPA Reg. No. 7594-6

DANGER

Keep Out Of Reach Of Children

Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food. Rinse empty container with water and discard it.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg white, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ACCEPTED WITH COMMENTS