


HOSPITAL BROAD BAND



USE DIRECTIONS:

Use at 1 oz. per gallon of water (1:128). Broad Band will provide 840 ppm phenols to this suggested use solution. This will provide disinfection against pathogenic organisms listed below, and consequently qualifies as a disinfectant for hospital use.

For heavily soiled areas, use 2 oz. per gallon of water.

Staphylococcus aureus

Sixty replicates on three preparations — also thirty replicates each on duplicate samples after sixty-day shelf-life study A.O.A.C. Method.

Salmonella Choleraesuis

Thirty replicates each on three preparations

Pseudomonas Aeruginosa

Thirty replicates each on three preparations

Manufactured By

**Sanitary** PRODUCTS CORP.
88 FARWELL STREET • WEST HAVEN, CONN. 06516
U.S.A.

CLEANER DISINFECTANT DEODORANT

ACTIVE INGREDIENTS

o-Benzyl-p-chlorophenol	5.00%
o Phenylphenol	4.25%
p-Tertiary amylphenol	1.25%
Tetrasodium ethylenediamine-tetraacetate	1.11%
Sodium Dodecylbenzenesulfonate	6.30%
Sodium xylene sulfonate	3.60%
Isopropanol	19.80%
INERT INGREDIENTS*	58.69%
	100.00%

*Contains cleaning and water softening agents. Broad Band is a concentrated, phosphate free, germicidal cleaner intended for cleaning and disinfection. It is designed for cleaning floors, walls, bed frames, furniture, and bathroom fixtures in a wiping, mopping, or scrubbing operation to wet surfaces thoroughly with the recommended use dilution.

It is free-rinsing and does not leave a film. Consequently, it is recommended for the cleaning of conductive flooring in operating rooms.

NET CONTENTS

EPA Reg. No. 7594-7

DANGER

**Keep Out Of Reach
Of Children**

Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food. Rinse empty container with water and discard it.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg white, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.