

KEM SAN

50

Disinfectant • Cleaner • Sanitizer
Fungicide • Deodorant

Active Ingredients:

Methyl dodecyl benzyl trimethyl ammonium chloride 80%
methyl dodecyl xylylene bis-trimethyl ammonium chloride) 20% 5.0%

Inert Ingredients 95.0%

DANGER: KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautionary handling instructions. Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

Manufactured by
CHEM LAB PRODUCTS INC.
2800 E. Coronado
Anaheim, California 92806

Net 55 Gallons

EPA Reg. No. 7616-24-AA
EPA Estab. No. 7616-CA-1

USE DIRECTIONS

FOR CLEANING: Lightly soiled areas - surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of KEM SAN 50 per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas: mop or brush soiled surface with a solution containing 2-3 ounces of KEM SAN 50 per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION: To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of KEM SAN 50 per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of KEM SAN 50 per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas where fungicidal action is required, mop area with a solution containing $\frac{3}{4}$ ounce per gallon of water or 3 ounces per 4 gallons of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION: To disinfect walls, floors, furniture and other hard surfaces in hospital and nursing homes, mop or brush surface to be treated with a solution containing $2\frac{1}{2}$ ounces of KEM SAN 50 per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION: To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing $\frac{3}{4}$ ounce of KEM SAN 50 per gallon of water. If surface is heavily soiled, use KEM SAN 50 at a rate of 2-3 ounces per gallon to ensure thorough cleaning. Sanitize cleaned surface with a rinse solution containing $\frac{1}{2}$ ounce of KEM SAN 50 per gallon of water. Those surfaces which will contact food must be rinsed with water of drinking quality before reuse.

DEODORIZATION: To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of KEM SAN 50 per gallon of water.

KEM SAN 50 is an effective antibacterial cleaner designed for institutional, industrial and institutional cleaning. When used as directed, KEM SAN 50 will disinfect hard surfaces such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where objectionable odors may develop. This product may be used as a general purpose antibacterial antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where cleaning and antimicrobial action is desired. See left panel for use direction.

DANGER: Keep Out of Reach of Children

Corrosive: Causes eye damage and skin irritation. Do not get in eyes or on skin or on clothing. Protect eyes and skin when handling. Handle if well diluted. Avoid contamination of food.

FIRST AID: In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eye: call a physician. Remove and wash contaminated clothing before reuse.

If swallowed: drink promptly a large quantity of milk, egg whites, saline solution or, if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may occur due to the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.

