

**STOCKADE
STOCKADE
STOCKADE
STOCKADE™**

Net Wt. 33 11/32 lbs.

**Bar-Fly
Fly Control Block**

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle when used as directed.

PESTICIDALLY ACTIVE DRUG INGREDIENT	
Phenothiazine	2.00%
PESTICIDALLY INERT INGREDIENTS	98.00%
TOTAL	100.00%

7627-11

GUARANTEED ANALYSIS

Crude protein, min.	1.000%
(This includes not more than 2.0% equivalent protein from non-protein nitrogen and not less than 8.0% protein from natural sources.)	
Crude fat, min.	1.000%
Crude fiber, max.	10.000%
Calcium (Ca), min.	2.000%
Calcium (Ca), max.	3.000%
Phosphorus (P), min.	1.500%
Iodine (I), min.	0.002%
Salt (NaCl), min.	22.500%
Salt (NaCl), max.	27.000%
Total Sugars (as invert), min.	20.000%
Vitamin A, USP units/lb.	60,000
Vitamin D3, USP units/lb.	12,500

FEED INGREDIENTS

Soybean meal, dried cane molasses, meat and bone meal, corn distillers dried grains with solubles, salt calcium carbonate, urea, monocalcium phosphate and dicalcium phosphate, vitamin A supplement, d-activated animal sterol, ethylenediamine dihydriodide, iron oxide, iron sulfate, zinc oxide, copper sulfate, manganous oxide, cobalt sulfate, animal fat with BHT, natural and artificial flavors, mineral oil, and cane molasses.

CAUTION!

KEEP OUT THE REACH OF CHILDREN

WARNING: Do not feed to lactating dairy animals. Harmful if swallowed. Avoid contact with skin and eyes. Wash thoroughly after handling and before eating or smoking. If in eyes, wash thoroughly with plenty of water for 15 minutes. If irritation persists, see a physician.

LIVESTOCK REMEDY - NOT FOR HUMAN USE!

Occasional sensitivity to phenothiazine has been reported and under certain conditions this drug is toxic to livestock. Animals that are sick, anemic, weak, emaciated or show signs of extreme constipation, should not be treated except upon the service of a veterinarian. Do not administer to pregnant animals in the last 4 weeks of pregnancy. Avoid exposure of recently treated animals to bright sunlight because of photosensitization. Do not use organophosphates or administer to animals showing symptoms of acute respiratory conditions.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

FEEDING DIRECTIONS

It is violation of Federal Law to use this product in a manner inconsistent with its labeling. Adult cattle must consume 3.5 ounces of Stockade Bar-Fly & Worm Control Block per head per day to obtain 2 grams phenothiazine. Younger cattle must consume 0.44 ounce per 100 lbs. body weight per head per day to receive 0.25 grams of phenothiazine intended. Continuous year round low-level phenothiazine treatment, in conjunction with good herd management practices, will control the specified parasites and will help prevent reinfection. This product kills horn fly and face fly maggots in manure, but does not control existing adult face flies. If necessary, spray cattle for maximum protection. Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult fly population.

STORAGE AND DISPOSAL

STORAGE: Do not store next to herbicides. Do not contaminate water or feed by storage or disposal. **CONTAINER DISPOSAL:** Completely empty container by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into mineral feeders. Then dispose of containers in a sanitary landfill or by incineration if allowed by State and Local authorities.

Manufactured by

EPA Reg. No. 7627-11 EPA Est. No. 7627-KS-01

HARVEST BRANDS, INC. Pittsburg, Kansas 66762



609



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

300/83092
17/2

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

Harvest Brands Inc.
P.O. Box 46
Pittsburg, KS 66762

Attn: Mr. Galen Williams

Dear Mr. Williams:

Subject: Update Labeling
Stockade Hi-Q Kil-A-Pest Fly Control Block
EPA Reg. No: 7627-15
Stockade Kil-A-Pest No. 12 Fly Control Block
EPA Reg. No: 7627-10
Stockade Bar-Fly Fly Control Block
EPA Reg. No: 7627-11
Appl. Date: October 20, 1982

The labeling for the product referred to above, submitted in connection with amended registrations under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, are acceptable, subject to the following comments, and stamped copies are enclosed for your records.

Prior to releasing the subject products for shipment under the amended labels, make the following revisions (for each product) and submit five copies of labeling, for each product, for our records:

1. Move the "KEEP OUT OF REACH OF CHILDREN" to appear immediately above the front panel signal word "CAUTION".
2. Delete the "!" following the front panel signal word.
3. Immediately preceding "WARNING: Do not feed to lactating dairy animals..." insert the following headings:

"PRECAUTIONARY STATEMENTS

HAZARDS TO ^{HUMANS} ~~HUMANS~~ AND DOMESTIC ANIMALS"

4. Delete the "WARNING" which appears at the beginning of this section: the signal word must be consistent throughout the labeling. "CAUTION" is the appropriate signal word to begin this section, however, since the "HAZARDS TO ^{HUMANS} ~~HUMANS~~..." section is on the front panel, the repetition of the signal word is unnecessary.
5. Revise the second sentence of the "STORAGE AND DISPOSAL" section to read:
"Do not contaminate water, food, or feed by storage or disposal."

(Note underline indicates added text and should not be reproduced on the label.)

-2-

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)