

**STOCKADE
STOCKADE
STOCKADE
STOCKADE®**

Net Wt. 50 lbs.

Fly Control Mixer

with Rabon® Oral Larvicide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle and house flies in the manure of treated swine.

PESTICIDALLY ACTIVE INGREDIENT

2-chloro-1-(2,4,5-trichlorophenyl)
vinyl dimethyl phosphate 0.40%*
INERT INGREDIENTS 99.60%**
Total 100.00%

*Each pound contains 1.81 grams Rabon.

RABON¹ is the registered trademark of Shell Oil Company.
U.S. Patent 3,102,842.

**Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein, minimum 4.50%
Crude Fat, minimum 3.25%
Crude Fiber, maximum 17.00%
Total Sugars
(as invert), minimum 38.00%

FEED INGREDIENTS

Dried cane molasses, animal fat (stabilized with BHT).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SWINE: All swine should be treated.

Mix 25 pounds Stockade Fly Control Mixer per ton of complete meal type feed and offer free-choice to hogs from weaning to market weight. This is equivalent to 22.7 milligrams of Rabon per pound of feed.

For sows, boars, and breeding gilts, mix 50 pounds of Stockade Fly Control Mixer per ton of meal type feed and offer 4-6 pounds of feed per animal per day. This is equivalent to 45.4 milligrams of Rabon per pound of feed.

CATTLE: For topdressing or use in mixed meal type feeds. Feed Stockade Fly Control Mixer to animals according to their live body weight by incorporating it into the daily ration or top dress in the grain, or roughage, as follows:

Body weight (lb.)	Stockade Rabon Mixer	
	(ounces)	(lb.)
400-1295	8	0.50
1295-1620	10	0.625
1620-1940	12	0.75

Stockade Fly Control Mixer may be fed up to slaughter and to lactating dairy cows. Start feeding this product early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

Stockade Fly Control Mixer prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle, but it is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases supplemental fly control measures may be needed in and around feed lots, calf pens and barns to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises. In order to achieve optimum fly control, Stockade Fly Control Mixer should be used in conjunction with other good sanitation and management practices.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Storage: Do not store next to herbicides. Do not contaminate water, food or feed by storage or disposal.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into mineral feeders, then dispose of bags in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CAUTION Must be fed and handled in accordance with labeling.



Manufactured by

HARVEST BRANDS, INC. Pittsburg, Kansas 66762

HARVEST®

074

EPA REG. NO. 7627-25
EPA EST. NO. 07627-KS-01

ACC

JAN 12 1987

7627-25