

7627-26

02/08/2007

111

STOCKADE Fly Control for Horses

with Rabon® Oral Larvicide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated horses.

ACTIVE INGREDIENT* Tetrachlorvinphos: (Z)-2-chloro-1 (2,4,5-trichlorophenyl) vinyl dimethyl phosphate 0.632%
OTHER INGREDIENTS** 99.368%

* contains 2.87 grams Rabon® Oral Larvicide per pound.
** refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein	minimum	8.00%
Crude Fat	minimum	0.50%
Crude Fiber	maximum	5.00%
Calcium (Ca)	minimum	7.50%
Calcium (Ca)	maximum	8.50%
Phosphorus (P)	minimum	5.00%
Salt (NaCl)	minimum	10.00%
Salt (NaCl)	maximum	12.00%
Magnesium (Mg)	minimum	0.50%
Copper (Cu)	minimum	25 ppm
Selenium (Se)	minimum	5 ppm
Zinc (Zn)	minimum	280 ppm
Iodine (I)	minimum	30 ppm
Vitamin A	minimum IU units/lb	30,000
Vitamin D3	minimum IU units/lb	7,500
Vitamin E	minimum IU units/lb	75

FEED INGREDIENTS: soybean meal, dried cane molasses, corn distillers dried grains with solubles, soybean mill feed, yeast culture, monocalcium phosphate, dicalcium phosphate, soft rock phosphate, salt, magnesium mica, calcium carbonate, iron oxide, calcium iodate, sodium selenite, zinc oxide, magnesium oxide, copper sulfate, cobalt carbonate, manganous oxide, ferrous carbonate, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, lignin sulfonate, cane molasses, natural and artificial flavors

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person. Do not induce vomiting unless told by a poison control center or doctor.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

IF ON EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Contains an organophosphate that inhibits cholinesterase. Have a product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-835-0306 for emergency medical treatment.

NOTE TO PHYSICIAN

This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 1-PAM is also antidotal and may be administered, but only in conjunction with atropine. Product may cause aspiration pneumonia. Probably mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with the eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters.

EPA Reg. No. 7627-26
EPA Est. No. 07627-KS-01

Manufactured by:

STOCKADE Brands, P.O. Box 46, Pittsburg, KS 66762

USER SAFETY REQUIREMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are butyl rubber, neoprene, and polyvinyl chloride (pvc). If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Loaders, mixers, applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

USE RESTRICTIONS:

This product is not to be used on horses destined for slaughter. This product contains tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the home are abdominal pain, lethargy sweating, tearing and excessive salivation. If these signs are seen in horses, consult your veterinarian immediately. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.

Allow free choice to horses. They should consume an average of 1.0 ounce of STOCKADE Fly Control for Horses per 300 pounds of weight per day. This will supply the recommended average daily intake of 70 mgs RABON Oral Larvicide per 100 pounds of body weight. Horses should not be starved for minerals before initiation of the Fly Control Block. Omit all salt from other feed after starting use of this block. Do not feed other free choice salt or mineral containing salt. One block should last a horse for approximately 130 days or 4 to 5 horses for 1 month. If consumption exceeds recommended level, reduce the number of feed locations or relocate. If consumption is below the recommended level, increase the number of feed locations. Begin feeding STOCKADE Fly Control Block for Horses early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall. This block prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated horses, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around horse lots and buildings to control adult house flies and stable flies which can breed not only in horse manure but in other decaying vegetable matter or silage.

STORAGE & DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in cool dry location in original container.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site in accordance with the label instructions or at an approved waste disposal facility. Open dumping is prohibited.
CONTAINER DISPOSAL: DO NOT reuse empty container. Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill, or incinerate, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

UPC CODE 088325006 **ACCEPTED** 0614
Net Wt. 33.33lb
15.12 kg

FEB - 8 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 7627-26