

50115

NET WEIGHT

Walnut Grove®

FLI-KIL 10 Mineral

With Rabon® Oral Larvicide



ACCEPTED

SEP 6 1984

Federal Insecticide,
and Rodenticide Act.
for the pesticide
under
EPA Reg. No. 7631-20

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle

ACTIVE INGREDIENTS:

2-chloro 1(2, 4, 5 trichlorophenyl)

vinyl dimethyl phosphate

INERT INGREDIENTS****BY WEIGHT:**

1.25%

98.75%

TOTAL

100.00%

Each Pound contains 5.625 g RABON

**Refers only to ingredients which are not larvicidal.

EPA REG. NO. 7631-20

EPA INT. NO.

7631-1A-1 7631-1A-3 7631-1A-5 7631-NB-1
7631-1A-2 7631-1A-4 7631-NB-1 7631-VT-1**FEED INGREDIENTS**

Wheat Middlings, Condensed Fermented Corn Extractives, Cane Molasses, Dried Molasses, Dicalcium Phosphate, Salt, Iron Oxide, Zinc Oxide, Manganese Oxide, Copper Sulfate, Cobalt Carbonate, Magnesium Oxide, Sodium Sulfate, Iron Sulfate, Calcium Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Mineral Oil, Ethoxyquin (a preservative).

GUARANTEED ANALYSIS

Calcium (Ca), maximum	12.00%
Calcium (Ca), minimum	10.00%
Phosphorus (P), minimum	10.00%
Salt (NaCl), maximum	11.00%
Salt (NaCl) minimum	9.00%
Magnesium (Mg), minimum	1.00%
Manganese (Mn), minimum	10%
Cobalt (Co), minimum	.001%
Copper (Cu), minimum	.01%
Iron (Fe), minimum	.15%
Vitamin A, minimum	300,000 U.S.P. Units Per Lb.
Vitamin D ₃ , minimum	45,000 U.S.P. Units Per Lb.
Vitamin E, minimum	150 IU/Per Lb.

KEEP OUT OF REACH OF CHILDREN**CAUTION**

SEE BACK OF LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured By

WALNUT GROVE PRODUCTS
Division of W. R. GRACE & CO.
General Offices
Atlantic, Iowa 50022



Walnut Grove® 4x4

FLI-KIL 10 Mineral

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

SPILLAGE: Store Walnut Grove 4x4 FLI-Kil 10 Mineral in a dry, well ventilated storage area away from other pesticides, animal feeds, herbicides, fertilizers, and manure. Store in original container with label intact to prevent loss of identity of product. Repair damaged bags promptly to prevent possible contamination of other products.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

EMPTYED BAGS: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

DIRECTIONS FOR USE: GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed flies: Feed to cattle. Cattle should consume an average of 1 oz. of the mineral per 500 lb. of body weight per day. This will supply the recommended average daily intake of 100 mg. of Rabon (carbicide) per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the Walnut Grove FLI-KIL 10 Mineral with Rabon program, and all salt from other feed after the program is started. Do not feed other feed containing salt or mineral containing salt in any form. Allow one mineral feeder for every 15 head, and place near watering and loafing areas. Put out no more than one feeder at a time, and protect from rain.

If consumption of Walnut Grove FLI-KIL 10 Mineral with Rabon is above 1 oz. per 500 lb. of body weight per day, reduce the number of feeding locations or relocate. If consumption is below 1 oz. per 500 lb. of body weight per day, increase the number of feeding locations. Start feeding Walnut Grove FLI-KIL 10 Mineral with Rabon early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Walnut Grove FLI-KIL 10 Mineral with Rabon may be fed up to slaughter and to lactating dairy cows.

Walnut Grove FLI-KIL 10 Mineral with Rabon prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing stable flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other unsightly matter or silage on the premises.

In order to achieve optimum fly control Walnut Grove FLI-KIL 10 Mineral with Rabon should be used in conjunction with other good management and sanitation practices.

Manufactured by

WALNUT GROVE PRODUCTS
Division of W. R. GRACE & CO.
General Offices
Atlantic, Iowa 50022