

GENERAL INFORMATION

Use TELONE II soil fumigant only as a **preplanting soil treatment** to control plant parasitic nematodes (root-knot, meadow lesion), citrus cyst formers (golden sugar beet, soybean), burrowing, ring spiral, sting, pin, stubby root, stylet dagger, also to control wireworms and garden centipedes (symphylans). Fumigate and to be planted to the crops listed below by applying TELONE II under the conditions and at the rates recommended under DIRECTIONS FOR USE, DOSAGE AND USE RECOMMENDATIONS, and USE PRECAUTIONS. Read the entire label before using TELONE II.

Vegetable Crops:

asparagus	cauliflower	horseradish	parsnips	shallots
beans	celery	kale	peas	spinach
beets	collards	kohlrabi	peppers	squash (summer)
blackeyed peas	corn	leeks	pimientos	squash (winter)
broccoli	cowpeas	lettuce	potatoes	sweet potatoes
brussels sprouts	cucumbers	melons	pumpkins	swiss chard
cabbage	egg plant	mustard greens	radishes	tomatoes
cantaloupe	endive	okra	rutabaga	turnips
carrots	garlic	onions	salsify	watermelons

Field Crops:

alfalfa	flax	oats	sorghum
barley	grasses	pasture grass	soybeans
birdsfoot trefoil	hops	peanuts	sugar beets
buckwheat	lespedeza	popcorn	sugarcane
clover	millet	rice	tobacco
corn	milo	rye	vetch
cotton	mint	safflower	wheat

Citrus Fruit Tree Planting Sites:

grapefruit	kumquats	lemons	limes	oranges	tangerines	tangelos
------------	----------	--------	-------	---------	------------	----------

Deciduous Fruit and Nut Tree Planting Sites:

almonds	dates	olives	plums
apples	figs	peaches	pomegranates
apricots	filberts	pears	prunes
cashews nuts	hazelnuts	pecans	quince
cherries	hickory nuts	persimmons	walnuts
chestnuts	nutrines	pineapple	

Bush and Vine Planting Sites:

blackberries	currants	huckleberries	youngberries
blueberries	dewberries	loganberries	
boysenberries	gooseberries	raspberries	
cranberries	grapes	strawberries	

Nursery Crops including floral plants, ornamentals, shrubs, and bushes, forest shade fruit and nut trees and vine and bramble fruits of all types.

**DOSAGE AND USE RECOMMENDATIONS
to Control Nematodes, Symphylans, and Wireworms:**

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage		
			Gallons Per Acre ¹	Fl. Oz./1000 Ft. Row Per Chisel	
Shallow Rooted Plants Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ⁶	Mineral	4.5 to 6	46 to 62	
		Muck or Peat	9 to 12	93 to 123	
	Overall (Broadcast)	Mineral	9 to 15 ²	26 to 44	
		Muck or Peat	24 ³ to 36	71 to 106	
Strawberries	Overall	Mineral	24 to 36	71 to 106	
Sugar Beets Root Knot Nematode Sugar Beet Nematode	Row (42")	Mineral	6	62	
	Overall		12 to 15	35 to 44	
	Row (42")		9	93	
	Overall		12 to 18	35 to 53	
Pineapple ⁴	Row	Mineral	24 to 36		
Citrus Florida ⁵	Overall	Mineral	36	106	
Nursery and Field Non-Fruit Trees Deciduous Fruit Trees Forest Trees	Overall Gallons Per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft	4 ft	5 ft	6 ft
	Sand	15	21	27	33

TELONE* II

SOIL FUMIGANT

A Clean, Clear, Non-Clogging Liquid for Preplant Treatment of Crop Lands to Control Plant Parasitic Nematodes and Certain Other Soil Pests

ACTIVE INGREDIENT:

1,3-Dichloropropene **92%**

INERT INGREDIENTS: **8%**

EPA Registration No. 7729-6

EPA Est. No. 7729 CA-3

Distributed by

JOHN TAYLOR FERTILIZERS CO.

AGRICULTURAL CHEMICAL

**Do Not Ship or Store with
Food, Feeds, or Clothing**

**PRECAUCION AL USUARIO: Si usted no lee
inglés, no use este producto hasta que la eti-
queta le haya sido explicada ampliamente.**

**TRANSLATION: (TO THE USER: If you cannot
read English, do not use this product until the
label has been fully explained to you.)**

DANGER

**KEEP OUT OF REACH OF CHILDREN
HAZARDOUS LIQUID AND VAPOR**

**MAY BE FATAL IF INHALED,
ABSORBED THROUGH SKIN OR SWALLOWED**

WHEN TO TREAT: Apply TELONE II (before October 15) is such as celery, tomatoes, a crop. **Note:** Treat muck soils (clay) soils only when they are wet when it is cold and/or wet.

SOIL PREPARATION: TELONE II is applied through the soil and the soil should be moist at about one-half inch. If undecomposed plant debris is present, TELONE II tillage deeper than 10 inches or "plow pan" occurs at plow to simultaneously apply the fumigant.

APPLICATION: TELONE II is applied with application equipment that will apply to the surface. For overall application, use chisel equipment to apply the fumigant up to twice the application depth. When the fumigant is applied, use chisel equipment to apply the fumigant to the soil where the crop is to be planted. Distribute about 1 1/2 times more fumigant per acre than the amount of fumigant required per acre of planting, avoid placing the fumigant in the center of the row.

SEALING: Immediately after chisel application, use a roller to compact the soil. Then compact it by floater wheel, by listing, or by bedder. Fumigating listed rows, seal the rows.

EXPOSURE PERIOD: After fumigation, the exposure period will be required.

AERATION AND PREPARATION: After the soil to aerate completely. TELONE II is no longer effective after 10 days for each 10 gallons of TELONE II and shrub planting sites. A 30-day period if heavy rains or low temperature occur. After row treatment, possible recontamination of the soil to the depth of the treatment of muck soils.

Important — Note Carefully: Do not use TELONE II in soils with high salt content. This is most common in either cold wet acid, or alkali soils. Tests made after fumigation of soils do not use fertilizers on the crop is well established.

Certain crops including cotton, corn, and soybeans should not be planted in the fumigated area. When using TELONE II in highly acid soils, ammonia toxicity. Certain non-vegetable crops such as carnation, avoid this possible effect. Experience indicates a definite effect on the crop.

Attention: To avoid re-planting

DOSE AND USE RECOMMENDATIONS
for Control of Nematodes, Symphytans, and Wireworms

For information on the use of this product, please refer to the label on the container. For a complete list of crops and pests controlled by this product, please refer to the label on the container.

Do Not Ship or Store with
Food, Feeds, or Clothing

PRECAUCION AL USUARIO. Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION TO THE USER. If you cannot read English, do not use the product until the label has been fully explained to you.

DANGER

KEEP OUT OF REACH OF CHILDREN
HAZARDOUS LIQUID AND VAPOR
MAY BE FATAL IF INHALED,
ABSORBED THROUGH SKIN OR SWALLOWED
CAUSES BURNS OF SKIN OR EYES
COMBUSTIBLE

Do Not Get in Eyes, on Skin, or on Clothing. Wear Eye and Skin Protection Necessary to Prevent Contact When Handling This Material. Wash Thoroughly after Handling and Before Eating and Smoking. Wash Contaminated Clothing and All Contaminated Shoes Thoroughly Before Reuse. DO NOT USE, POUR, SPIEL, OR STORE NEAR HEAT OR OPEN FLAME. Use Only with adequate Ventilation. Do Not Breathe Vapor.

JOHN TAYLOR FERTILIZERS CO

TELONE II

SOIL FUMIGANT

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions. Do not use on crops listed unless you have read the label carefully and understand the directions.

ACTIVE INGREDIENT
1,3-Dichloropropene 92%

INERT INGREDIENTS 8%

Net Weight 100 lbs. (45.4 kg.)
Gross Weight 110 lbs. (50.0 kg.)

Distributed by
JOHN TAYLOR FERTILIZERS CO.

AGRICULTURAL CHEMICAL

Do Not Ship or Store with
Food, Feeds, or Clothing

PRECAUCION AL USUARIO: No usee este producto en los campos que no se listan en el rótulo. No usee este producto en los campos que no se listan en el rótulo. No usee este producto en los campos que no se listan en el rótulo.

TRANSLATION (TO THE USER): If you cannot read English, do not use this product until the label has been fully explained to you.

DANGER

KEEP OUT OF REACH OF CHILDREN.
DO NOT USE IF CONTAINER IS OPENED.
DO NOT USE IF CONTAINER IS OPENED.
DO NOT USE IF CONTAINER IS OPENED.

DIRECTIONS FOR USE

WHEN TO TREAT

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

APPLICABLE CROPS

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

APPLICATION

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

EXPOSURE PERIOD

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

PREPARATION AND REPAIRATION OF

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

DIRECTIONS FOR USE

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

APPLICABLE CROPS

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

EXPOSURE PERIOD

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

PREPARATION AND REPAIRATION OF

Use only on crops listed on label. Do not use on crops not listed. Do not use on crops listed unless you have read the label carefully and understand the directions.

