## LABELS FOR ROCCAL-D PACKETS

FRONT PANEL

# ROCCAL®D

VIRUCIDE\*

DISINFECTANT - DETERGENT - DEODORANT

#### Dilute with water as directed

Rinse empty container thoroughly with water and discard it.

"Activity demonstrated against Canine Distemper virus, Canine Herpes virus, and Pseudorabies virus at dilutions recommended for treatment of hard inanimate surfaces.

# DANGER: Keep out of reach of children.

Corrosive to tissues. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling the concentrate. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

1/3 FL. OZ. (10 ML.)

## REAR PANEL

Roccal-D is a pleasantly scented, colored solution of high antibacterial and antiviral potency for treatment of hard inanimate surfaces.

Active Ingredients

Alkyl\* dimethyl benzyl ammonium chlorides 20 00%

\*(67% C ... 25% C ... 7° ... C ... 1° ... C ... c ... and C ...)

Monoethanolamine ethylenediamine tetraacetate 1 75° ...
Essilatiation 2 52° ...

Inert Ingredients ... 75 73° ...

#### DIRECTIONS FOR USE

FOR GENERAL DISINFECTION OF HARD SURFACES, such as floors, walls, operating tables, cages and furniture in animal hospitals, kennels, animal quarters and laboratories: Clean thoroughly and rinse all surface areas before disinfecting. Then apply Roccal-D dilution 1:400 by mop, sponge, or spray and let air dry. For surfaces difficult to clean and heavily contaminated areas, use Roccal-D dilution 1:200.

FOR DISINFECTION OF EATING AND DRINKING UTENSILS for animals not considered as food sources for human consumption: Scrape and prewash with warm water, then soak in Roccal-D dilution 1:400 for at least 10 minutes, and rinse with potable water.

### PREPARATION OF USE DILUTIONS

**DILUTION 1:400** Add 10 ml. (1 packet) of Roccal-D to 1 gallon of water.

**DILUTION 1:200** Add 20 ml. (2 packets) of Roccal-D to 1 gallon of water.

Do not mix Roccal-D with soap or anionic detergents

EPA Reg No 7738-3-AA EPA EST 944-PA-1



Veterinary Department
Winthrop Laboratories Division of Sterling Drug Inc
New York, N.Y. 10016

**RW** 24

Made in U.S.A.

## LABELS FOR ROCCAL-D PACKETS

FRONT PANEL

# ROCCAL®D

**VIRUCIDE\*** 

DISINFECTANT - DETERGENT - DEODORANT

#### Dilute with water as directed

Mode energy container theroughly with water and discard it.

\*Activity demonstrated against Canine Distancer virus, Canine Horpes virus, and Pacuderables virus at dilutions recommended for transferent of hard irranimate surfaces.

# DANGER: Keep out of reach of children.

Corrosive to tissues. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling the concentrate. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

MOTE TO PHYSICIAM: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulator, shock, respiratory depression, and convulsion may be needed.

1/3 FL OZ. (10 ML)

## REAR PANEL

Roccal-D is a pleasantly scented, colored solution of high antibacterial and antiviral potency for treatment of hard inanimate surfaces.

#### DIRECTIONS FOR USE

FOR GENERAL DISINFECTION OF HARD SUR-FACES, such as floors, walls, operating tables, cages and furniture in animal hospitals, kennels, animal quarters and laboratories: Clean thoroughly and rinse all surface areas before disinfecting. Then apply Roccal-D dilution 1:400 by mop, sponge, or spray and let air dry. For surfaces difficult to clean and heavily contaminated areas, use Roccal-D dilution 1:200.

FOR DISINFECTION OF EATING AND DRINKING UTENSILS for animals not considered as food sources for human consumption: Scrape and prewash with warm water, then soak in Roccal-D dilution 1:400 for at least 10 minutes, and rinse with potable water.

### PREPARATION OF USE DILUTIONS

DILUTION 1:400 Add 10 ml. (1 packet) of Roccal-D to 1 gallon of water.

**DILUTION 1:200** Add 20 ml. (2 packets) of Roccal-D to 1 gallon of water.

- Do not mix Roccal-D with soap or anionic detergents.

EPA Reg. No. 7738-3-AA EPA EST 944-PA-1



Veterinary Department
Winthrop Laboratories Division of Sterling Drug Inc.
New York, N.Y. 10016

RW 24

Made in U.S.A.