


**ACCEPTED**  
 MAY 8 1976  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 7762-12

EPA  
 507

Vertical text on the left edge, partially obscured and difficult to read.



**COMBATO PROMIX**  
 NET WT. 25 LBS.


(WITH 7.76% CYALUMINIDE 7.76%)

TO PREVENT THE DEVELOPMENT OF HOUSEFLIES, FACE FLIES, HOUSE  
 FLIES AND STABLE FLIES IN THE BARN OF TREATED CATTLE.

ACTIVE INGREDIENTS	BY WEIGHT
Chloro-1-(2,4,6-trichlorophenyl) vinyl	7.76%
Dimethyl phthalate	92.24%
<b>TOTAL</b>	<b>100.0%</b>

Small text at the bottom of the page, likely a registration or disclaimer notice.

25 LBS.



(WITH 7.76% ORAL LARVICIDE 7.76%)

TO PREVENT THE DEVELOPMENT OF HORSE FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANNING OF TREATED CATTLE.

ACTIVE INGREDIENT	BY WEIGHT
3-chloro-1-(2,4,6-trichlorophenyl) Vinyl	7.76%
Inert ingredients	92.24%
<b>TOTAL</b>	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN**

**PRECAUTIONARY STATEMENTS**

**HAZARDOUS TO HUMANS**

Harmful if swallowed. Avoid contact with eyes. If contact occurs, wash thoroughly with soap and water. If getting on skin, wash with plenty of water for 15 minutes. If on clothes, wash immediately.

**ENVIRONMENTAL PRECAUTIONS**

This product should not be used in areas where it may contaminate water by cleaning of equipment or disposal.

NET WEIGHT 25 LBS. (11.34 kg) EPA REG. NO. 7282-12

NET CONTENTS 25 LBS. (11.34 kg) EPA REG. NO.

PREPARED BY DEWIS COMPANY, P.O. BOX 1000, CALIFORNIA 94500

REGISTERED TRADEMARK (1972)

PREMIER FLY CONTROL

# DIRECTIONS FOR USE

## GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LESLIE FLY CONTROL "R" PREMIX will mix uniformly in cattle feeds when standard mixing procedures are followed. The usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution of LESLIE FLY CONTROL "R" PREMIX throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if LESLIE FLY CONTROL "R" PREMIX is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer feeder truck) may be used to prepare formulated feeds.

Feeds prepared with LESLIE FLY CONTROL "R" PREMIX should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, LESLIE FLY CONTROL "R" PREMIX should not be mixed in liquid feed supplements.

When fed to cattle, LESLIE FLY CONTROL "R" PREMIX passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, LESLIE FLY CONTROL "R" PREMIX helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

LESLIE FLY CONTROL "R" PREMIX prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, LESLIE FLY CONTROL "R" PREMIX should be used in conjunction with other good management and sanitation practices.

### In a Complete Ration—No Other Roughage Fed:

LESLIE FLY CONTROL "R" PREMIX can be used to prepare rations that contain 26.4 mg of RABON per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of LESLIE FLY CONTROL "R" PREMIX per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

### In a Concentrate Feed—Roughage Fed Separately:

7762-12

Cameraman's Note

Poor Copy

## Cameraman's Note

Poor Copy

during a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, LESLIE FLY CONTROL "R" PREMIX should be used in conjunction with other good management and sanitation practices.

### **In a Complete Ration—No Other Roughage Fed:**

LESLIE FLY CONTROL "R" PREMIX can be used to prepare rations that contain 26.4 mg of RABON per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of LESLIE FLY CONTROL "R" PREMIX per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

### **In a Concentrate Feed—Roughage Fed Separately:**

LESLIE FLY CONTROL "R" PREMIX can be used to prepare concentrate feeds that will provide 792 mg of RABON per animal per day.

To prepare a larvicidal concentrate feed, mix LESLIE FLY CONTROL "R" PREMIX according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

Pounds of LESLIE FLY CONTROL "R" PREMIX per ton of Concentrate	MIXING DIRECTIONS		Pounds of Concentrate Consumed per Animal per day
	RABON in the Concentrate mg/lb	%	
90.0	1584	0.35	.5
45.0	792	0.18	1.0
30.0	528	0.12	1.5
22.5	396	0.087	2.0
9.0	159	0.035	5.0
4.5	79	0.018	10.0
3.0	53	0.012	15.0
2.3	39	0.0087	20.0
1.8	32	0.0059	25.0
1.5	26	0.0059	30.0

Feed the appropriate larvicidal concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200-1700 pounds, increase the amount of concentrate fed to 1½ times that recommended.

Start feeding LESLIE'S RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Rations containing LESLIE'S RABON Oral Larvicide may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

## **STORAGE AND DISPOSAL**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.