


ACCEPTED  
7702 13



DATE  
TIME

LESIE  
SALT  
CO.  
NEWARK  
CALIFORNIA  
94566

# Leslie

## FLY CONTROL NO. 11-R

(WITH FABEN<sup>®</sup> ORAL LARVICIDE)

TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE  
FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate .....	1.8%
INERT INGREDIENTS* .....	98.2%
TOTAL .....	100.0%

\*Not all of the ingredients listed are active.

GUARANTEED ANALYSIS	
CALCIUM (Ca) .....	Minimum 14.00%
PHOSPHORUS (P) .....	Minimum 4.50%
SALT .....	Minimum 30.00%
SALT .....	Maximum 34.00%

INGREDIENTS Salt, Limestone, Dicalcium Phosphate, Fine Meal, Dried Molasses, Cane Molasses, Condensed Extract of Lactic Acid Fermentation Product, Manganese Sulfate, Zinc Oxide, Iron Oxide, Selenium, Iron Carbonate, Copper Oxide, Calcium Iodate, Borax, Gel, Cobalt Carbonate.

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WT. 50 LBS. EPA REG. NO. 7702-13  
EPA IDENT. NO.

PRODUCED BY LESLIE SALT CO. NEWARK, CALIFORNIA 94566  
INFO. BY LESLIE SALT CO. + NEWARK, CALIFORNIA 94566  
Phone: (415) 737-1220

DATE  
TIME

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

7762-13

## DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of  $\frac{1}{4}$  lb. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg RABON Oral Larvicide per hundred pounds body weight. Animals should not be starved for minerals before initiation of the RABON Cattle Mineral Mix program. Do NOT OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 15-20 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Leslie Fly Control No. 11-R is above  $\frac{1}{4}$  lb. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below  $\frac{1}{4}$  lb. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Leslie Fly Control No. 11-R early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Leslie Fly Control No. 11-R may be fed up to slaughter and to lactating dairy cows.

Leslie Fly Control No. 11-R prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Leslie Fly Control No. 11-R should be used in conjunction with other good management and sanitation practices.

## STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.