

NO. 11-R

WITH RABON® ORAL LARVICIDE

TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE.

ACCEPTED

1:AY 9 5 18/10

Unit the form I Insertifies, Funciale, et al. of all Act. du driet und, . . o prabeide required under

EPA Res N. 7742

ACTIVE INGREDIENTS

BY WEIGHT

2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate

1.8%

INERT* INGREDIENTS

98.2% TOTAL 100.0%

*Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

CALCIUM (Ca) CALCIUM (Ca) PHOSPHORUS (P) Minimum 14.00% Maximum 16.00% Minimum 4.50% SALT 30.00% Minimum SALT 34.00% Maximum

INGREDIENTS: Salt, Limestone, Dicalcium Phosphate, Cane Molasses, Condensed Extracted Glutamic Acid Fermentation Product, Bone Meal, Dried Molasses, Ammonium Lignin Sulfonate, Manganese Sulfate, Zinc Oxide, Iron Oxide, Sulfur, Iron Carbonate, Copper Oxide, Calcium Iodate, Mineral Oil, Cobalt Carbonate, Fenngreek, Anise Oil.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

NET WT. 40 LBS. EPA REG. NO. 7762-14 EPA EST. NO.

RABON' IS THE REGISTERED TRADEMARK OF SHELL CHEMICAL COMPANY, U.S. PATENT 3,102,842

MFG. BY LESLIE SALT CO. * NEWARK, CALIF. 94560

Physe: (415) 797-1820

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

7766.14

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

K.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 15-20 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of ¼ lb. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the RABON cattle Mineral Block program but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain.

If consumption of Leslie Fly Control Block No. 11-R is above ¼ ib. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below ¼ lb. of the block per 100 lbs. of body weight per month, increase > number of feeding locations. Start feeding Leslie Fly Control Block No. 11-R early in the spring before flies begin to appear, and continue feeding throughout the summer and the fall.

Leslie Fly Control Block No. 11-R may be fed up to slaughter and to lactating dairy cows.

Leslie Fly Control Block No. 11-R prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program, during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental by control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manufe but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Leslie Fly Control Block No. 11-R should be used in conjunction with other good management and sanitation practices.

7762-14

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER OR FOOD by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.