


**SYSTEMIC DANDELION & CHICKWEED KILLER**

**COVERS 5,000 SQ. FT.**

# SYSTEMIC DANDELION & CHICKWEED KILLER



**Controls Most Broadleaved and Vining-type Weeds**

- (1) Combines the benefits of 2, 4-D and Silvex, two of the most effective Weed Killers available.
- (2) Kills Dandelions, Chickweed, Ground Ivy and most other broadleaved and vining-type lawn weeds such as:

Buckhorn	Docks	Hent:	Pennywort
Chickweed	Dandelion	Knotweed	Plantain
Cinquefoil	Fieldbindweed	Annual Morningglory	Purslane
Clovers	Ground Ivy	Oxalis	Thistles


**Active Ingredients**

2,4-Dichlorophenoxyacetic acid (as its dimethylamine salt)	0.77%
2,4,5-Trichlorophenoxyacetic acid (as its dimethylamine salt)	0.60%
2,4-Dichlorophenoxyacetic acid (as its dimethylamine salt)	10.40%
2,4,5-Trichlorophenoxyacetic acid (as its dimethylamine salt)	10.40%
2,4,5-Trichlorophenoxyacetic acid (as its dimethylamine salt)	0.60%
2,4,5-Trichlorophenoxyacetic acid (as its dimethylamine salt)	0.60%

**Caution: Keep Out of Reach of Children.** Avoid contact with eyes, skin and clothing. Avoid breathing dust. Avoid application to garden plants, shrubs, and trees. Do not use in lawns or other areas where children play. Do not use on lawns, shrubs, trees, or other plants. Do not use on lawns, shrubs, trees, or other plants.

© 1970 HERITAGE HOUSE, Dept. of Research, Division of Agriculture, a unit of Heritance Research Corporation—HERITAGE, 600 0013

SPEC. NO. HH-20-A-2  
G-37158



K9

**HERITAGE HOUSE**


**SYSTEMIC DANDELION & CHICKWEED KILLER**

**COVERS 5,000 SQ. FT.**

# DANDELION & CHICKWEED KILLER

**Covers 5000 Sq. Ft.**

**Controls Most Broadleaved and Vining-type Lawn Weeds Including Dandelion, Chickweed and Clover.**



**Directions**

**When To Apply:** Use on lawn weeds that are actively growing and when temperature reaches 50 degrees or above. Control is slower if weather is cool. For optimum results apply when moisture is on grass.

**How To Apply:** Apply over entire lawn area. Use a spreader set for an application rate of 4 Lbs. per 1,000 Sq. Ft. Shut off spreader as you stop or turn around. Rain or sprinkling, mowing or heavy traffic within 24 hours reduces effectiveness and a repeat treatment may be needed—wait at least one week before planting new grass for two weeks. Do not apply to new plantings until after new grass has been mowed at least twice. Grass may become discolored temporarily in periods of high heat or humidity, but with watering will make a fairly rapid recovery.

**Spreaders**

Spreader Model	Normal Rate	Grassroots
Heritage Wheel Drop Spreader Model #1002	4	5
Heritage Broadcast Model D 2712	4	5
Cyclone Model #B1	4 1/2	5 1/2
New Beauty 24" Drop Model 2455	4	5
Scout 24" Drop Model 753	4	5
Scout 18" Drop Model 353	4	5
Scout 12" Drop Model 15219202	4	5
Green 20" Drop Model #205	4	5
Heritage Broadcast Model #2711	4	5

**Spreader Settings Caution:**

Note: The settings for spreaders other than Heritage House are only approximate. Age and condition of spreader will affect application rate. Observe the application rate on a test area before applying to the lawn. If the application rate is too high, the product will be applied unevenly. If the application rate is too low, the product will not be applied evenly. Do not over-apply. The recommended rate which can be used on any type of grass.


Caution: Do not use when there is a hazard of drift to other plants. Do not apply to lawns, shrubs, trees, or other plants. Do not use on lawns, shrubs, trees, or other plants.

Warning: Do not use on lawns, shrubs, trees, or other plants. Do not use on lawns, shrubs, trees, or other plants. Do not use on lawns, shrubs, trees, or other plants.

**ACCEPTED**  
*2-11-70*  
**UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 7767-18**

**HERITAGE HOUSE**


# SYSTEMIC DANDELION CHICKWEED KILLER



type Weeds  
st effective Weed

ner broadleaved

Pennywort  
Plantain  
Purslane  
Thistles



**What Corncob Chemistry Means**

The carrier in this product is organic—dry corn cob granules saturated with weed control solutions. Each corn cob granule has been impregnated with the proper ingredients. Result—a uniform, lightweight product that releases the ingredients evenly as the Corn Cob granule decomposes due to moisture and temperature.

Because of its organic nature, the corn cob granule is subject to decay when in contact with the soil and decomposes into Humus—a desirable benefit.

## Directions

### When To Apply

Use anytime weeds are actively growing and when temperature reaches 50 degrees or above. Control is slower if weather is cool. For optimum results apply when moisture is on grass.

### How To Apply

Apply over entire lawn area. Use a spreader set for an application rate of 4 Lbs. per 1,000 Sq. Ft. Shut off spreader as you stop or turn around. Rain or sprinkling, mowing or heavy traffic within 24 hours reduces effectiveness, and a repeat treatment may be needed—wait at least one week. Delay planting new grass for two weeks.

Do not apply to new plantings until after new grass has been mowed at least twice.

Grass may become discolored temporarily in periods of high heat or humidity, but with watering will make a fairly rapid recovery.

## Spreaders

	Normal Rate	Crisscross
HH (White) 20" Drop, Spreader Model #1662	6	5
HH Broadcast, Model D-2712	4	3
Cyclone Model #B1	4½	3½
Lawn Beauty 24" Drop, Model 2455	7	5½
Scott's 24" Drop, Model 75-3	5½	4½
Scott's 18" Drop, Model 35-3	6½	5½
Sears Craftsman 24" Drop, Model 45219202	4½	3½
Greenfield 20" Drop, Model #205	7	5½
True Temper Broadcast, Model #D2711	4	3

### Spreader Settings Caution:

**Note:** The settings for spreaders other than Heritage House are only approximates. Age and condition of spreader will affect application. User should carefully observe rate of application at beginning of spread and make adjustment if necessary. Our products have been tested on new spreaders only. Over application of this product can produce serious turf injury. Do not open spreader while in stationary position or while making turns, as you will be applying seven to ten times the recommended rate, which can cause severe burning to grass.

**Caution:** Use only when there is no hazard of drift. May injure bentgrass, St. Augustine, D.chondra, carpetgrass, clover and grass seedlings in newly seeded ground. Vapors from this product may injure susceptible plants in the immediate vicinity.

**Warning:** Do not apply directly to or otherwise permit it to contact vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other desirable plants which are sensitive to 2, 4-D and similar materials. Excess amounts of 2, 4-D Acid in the soil may temporarily inhibit seed germination or plant growth. Do not use on crops that will be utilized for food or forage purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water used for domestic or irrigational purposes.



HERITAGE HOUSE

COVERS 5,000 SQ. FT.  
\$4.45

HERITAGE HOUSE


HERITAGE HOUSE

TO OPEN: TAP END CAP  
AND PULL OUT COVER TUBE

HERITAGE HOUSE

SYSTEMIC DANDELION & CHICKWEED KILLER

# SYSTEMIC DANDELION & CHICKWEED KILLER



Controls Most Broad leaved and living type Weeds

Active Ingredients

Caution: Keep Out of Reach of Children

HERITAGE HOUSE


SYSTEMIC DANDELION & CHICKWEED KILLER

COVERS 5,000 SQ. FT.

ACCEPTED  
2-11-70  
7764-18

# DANDELION & CHICKWEED KILLER

Covers 5000 Sq. Ft.



Directions

HERITAGE HOUSE

