

DIRECTIONS FOR USE

Put about a cup of water in a watering can or other container. Add one fluid ounce (2 tablespoonsful) of SLUGIT. Then add one gallon water and stir thoroughly. Apply this quantity of solution by spraying through a coarse nozzle or with a water can to an area of 100 square feet. Make sure the entire area, including plants, is covered. A second application about 10 days later is recommended.

SLUGIT is not known to be harmful to plants when used as directed, but as an added precaution, a test should be made before applying it to valuable specimens of unusual plants. SLUGIT is harmless to earthworms and beneficial insects when used as directed. For best results, apply when slugs are active and when rain is not immediately expected. SLUGIT remains effective for about one week or until it is washed off by rain or other means.

Do not take internally. If swallowed, give an emetic of a tablespoonful of salt in a glass of warm water.

Do not use on or around food plants.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

THIS PRODUCT IS TOXIC TO WILDLIFE. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED.

**SHAKE CONTAINER WELL BEFORE USE
PROTECT FROM FROST
KEEP SOLUTION WELL STIRRED**

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury, whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

'Murphy' is a registered trade mark.

USDA Registration No. 7782-1

Made in England by



Murphy Chemical Ltd

Wheathampstead, St Albans, Hertfordshire, England

USA0285A, 1/72, T

**Murphy
Slugit
Liquid**

**Murphy
Slugit
Liquid**

**Active ingredient: Metaldehyde 20%
Inert ingredient: 80%**

GENERAL INFORMATION

BEFORE USING THIS PRODUCT - READ THE LABEL CAREFULLY

PAY PARTICULAR ATTENTION TO THE PRECAUTIONS AND WARNING STATEMENTS

If conditions as set forth on this label are not acceptable, return the unopened container and contents at once for full credit or refund of full purchase price.

The suggestions and Directions for Use of this product are based on information we believe to be reliable and appropriate. They are made in good faith, but without guarantee.

Severe crop damage or other unintended results may follow the use of this product because of unfavorable growing conditions; unfavorable weather; presence of other products or materials in or on the soil or crop being treated; or because of the method of use or application.

Specific instructions to cover all contingencies surrounding the use of this product are beyond the scope or space on a product label, therefore the exercise of care and good judgement in the use of this material is urged, particularly when unusual weather, soil conditions, or weed or crop growth are involved.

Consult State Agricultural Experiment Station and Extension Service Specialists for additional information concerning unusual local problems involving the use of this material.

- - - - -

This product is extremely toxic and should be handled only by experienced personnel using approved safety equipment. It is formulated for mixing with water before application. Apply with conventional hydraulic sprayers, low volume ground machines or airplane sprayers.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

Do not apply during bloom when honey bees and other pollinating insects are active.

Workers entering treated fields within 24 hours after application should wear protective clothing.