

RBC - RBC - RBC - RBC

# RED BARN MALATHION 5P

RBC - RBC - RBC

## GENERAL DIRECTIONS

Before using this material be sure that spray tank, supply lines, hose and nozzle are in good condition; that hose is sound and securely fastened; that nozzles are adjusted and the machine is properly calibrated to deliver the desired amount of spray.

When using with backpack sprayer continue agitation while spraying and agitate thoroughly before using.

For best results mix the required amount of this material with sufficient water to give uniform coverage. Time applications in accordance with state and federal recommendations. Suggested rates of application are given for specific crops listed on this label.

## OBSERVE THESE PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams, and ponds.

Do not apply during bloom when honey bees and other pollinating insects are active.

## DIRECTIONS FOR USE

**ON ALFALFA** — For control of spotted alfalfa aphids use 1 to 1 1/2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary to maintain control.

For control of aphids, grasshoppers, leafhoppers, spider mites and spittle bug use 1 1/2 to 2 pints per acre.

Do not apply during full bloom or within 7 days of harvest.

**ON BEETS** — For control of aphids use 1 1/2 to 2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 7 days of harvest if tops are to be used.

**ON COLE CROPS** (Broccoli, Brussels Sprouts, Cabbage, Cauliflower) — For control of aphids, imported cabbage worm, cabbage looper use 1 to 2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 7 days of harvest on Brussels Sprouts, Cabbage, Cauliflower or within 3 days of harvest on Broccoli.

**ON CELERY** — For control of aphids and spider mites use 1 1/2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 7 days of harvest.

## EMULSIFIABLE CONCENTRATE

Contains 5 Lbs. Premium Grade Malathion Per Gallon

### ACTIVE INGREDIENTS:

Malathion\* 56%  
Xylene 34%

### INERT INGREDIENTS

TOTAL 100%

\*O, O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

## CAUTION — Keep Out of the Reach of Children

Harmful if Swallowed

Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Keep container closed and away from heat, sparks and open flame.

USDA Reg. No. 7794-47

**NOTICE:** The information contained herein is, to our best knowledge, true and accurate. Because conditions of use which are of critical importance are beyond our control, we make no warranty or representation, express or implied, except that all products conform to the chemical description on the label. No agent of this company is authorized to make any warranty or representation concerning our products. Always follow directions and carefully observe all precautions on the label. Products used contrary to directions may cause serious plant or personal injury. We are not responsible for chemicals after they leave our possession.

CONTENTS \_\_\_\_\_ GALLONS



FERTILIZERS • CHEMICALS

RED BARN CHEMICALS, INC.

SHREVEPORT, LOUISIANA

**ON LETTUCE** — For control of aphids and leafhoppers use 2 pints per acre. For cabbage loopers use 3 pints per acre. Mix in sufficient water for adequate coverage. Repeat as necessary.

Do not apply to head lettuce within 7 days of harvest or to leaf lettuce within 14 days.

**ON PEAS** — For control of pea aphid and pea weevil use 1 1/2 to 2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 7 days of harvest if vines are to be used for silage or within 3 days if vines are not to be fed.

**ON PECANS** — For control of aphids and spider mites use 1 to 2 gallons of water. Make full coverage applications of spray as necessary.

**ON SPINACH** — For control of aphids use 2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 7 days of harvest.

**ON TOMATO** — For control of aphids and spider mites use 1 to 2 pints per acre in sufficient water to secure adequate coverage. Repeat as necessary.

Do not apply within 1 day of harvest.

For control of insects such as the confused flour beetle, fat grain lesser grain beetle, red flour beetle, saw-toothed grain beetle, weevil, rice weevil, and Indian meal moth on stored grains — wheat, rye, barley, corn, grain sorghum, peanuts, rice, field & garden.

For each 1000 bushels to be treated mix 1 pint of Malathion 5P in 10 gallons of water. Apply uniformly to the grain as it is being dumped into final storage.

To protect against Indian meal moth spray grain at the rate of 1/2 Malathion 5P per 1000 square feet of exposed surface area (annular) after being stored. Repeat treatment if necessary.

Before storage of grain clean truck beds, tank cars, elevators, the bins, and remove all nesting and debris.

Mix 1 gallon of Malathion 5P in 25 gallons of water & apply as a residual to walls, floors, truck beds, box cars, ships, bins as a residual spray.

RBC - RBC - RBC - RBC

MALATHION 5P

7794-47

