



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Read these entire Directions and Conditions of Sale before using BORDERLAND BLACK REPELLENT.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. FAILURE TO FOLLOW THESE DIRECTIONS OR THE OCCURRENCE OF WEATHER CONDITIONS THAT ARE EXTRAORDINARY OR UNUSUAL FOR THE USE AREA MAY RESULT IN UNSATISFACTORY PERFORMANCE OF THE PRODUCT.

BORDERLAND PRODUCTS, INC. AND THE SELLER OFFER, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE AFOREMENTIONED CONDITIONS WHICH ARE BEYOND THE CONTROL OF BORDERLAND PRODUCTS, INC. AND THEREFORE THE RESPONSIBILITY OF THE BUYER.

BORDERLAND BLACK Repellent is effective in preventing bird feeding damage to newly planted corn (including field, sweet, and popcorn) in all States east of the Mississippi River only. Blackbirds are repelled from treated fields before they can cause significant loss of seeds or seedlings. Birds feeding on treated seed may become paralyzed, however, the majority recover within a short period of time.

Apply **BORDERLAND BLACK Repellent** at planting at a rate of 14.3 oz. per 100 pounds of seed corn (8 oz. per bushel). Place the appropriate amount of **BORDERLAND BLACK Repellent** in the filled seed hopper. Using a long stick or rod, stir the hopper contents to thoroughly mix the bird repellent with the seed. Check equipment frequently to assure proper operation.

Under cool, wet growing conditions, this product may be phytotoxic to corn.

STORAGE AND DISPOSAL

1. **PROHIBITIONS.** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

2. **PESTICIDE DISPOSAL.** Pesticide or its waste that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

3. **CONTAINER DISPOSAL.** Dispose of in a landfill approved for pesticide containers or bury in a safe place.

4. **GENERAL.** Consult federal, state, or local disposal authorities for approved alternative procedures such as limited open burning.

Sold by

BORDERLAND PRODUCTS, INC.

P.O. Box 366, Buffalo, N.Y. 14240

STATEMENTS OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
If on skin, remove contaminated clothing and wash skin thoroughly with soap and warm water. If eyes are contaminated, wash eyes with water for at least 15 minutes.
Treated seed must not be used for feed, food, or forage. Use only according to label directions. Applications at rates above those shown may result in reduced crop residues.
This pesticide is toxic to fish and other wildlife. Birds feeding on treated corn may be killed. Keep out of lakes, ponds, and streams. Dispose of excess treated seeds or wastes by burial away from water supplies. Do not allow treated seeds to be exposed on soil surface. Apply only as specified on the label.
MESUROL is a Reg. TM of the Parent Company of Fermentation
Bayer GmbH, Leverkusen.