





Squire Applegate
TM



Squire Applegate
TM



5% SEVIN
DUST

2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

To control certain white grubs (larvae of European Chafer and Japanese beetle): Use two to four times the amounts of Dursban 2 EC shown in the turf table, i.e. 3 to 6 fl. oz. per 1000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July or August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

Pests of Ornamentals: Use Dursban 2 EC to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed) and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute Dursban 2 EC with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. Note—Do not use on azaleas, camelias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

DOSAGE and DILUTIONS for USE on ORNAMENTAL PLANTS

Pests Controlled	Amount of Dursban 2 EC in Water to Make		
	3 gallons	10 gallons	100 gallons
Bagworms			
Grasshoppers			
Mites	0.5 fl. oz.	1.6 fl. oz.	1 pint*
Spittlebugs			
Whiteflies			
Cutworms			
Mealybugs			
Leafhoppers	1 fl. oz.	3.2 fl. oz.	1 quart**
Thrips (exposed)			

*Equivalent to 4 ounces of chlorpyrifos per 100 gallons.

**Equivalent to 8 ounces of chlorpyrifos per 100 gallons.

Note: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

Warning (Lawn and Ornamental Plant Pests): Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water. Fish may be killed if their waters are contaminated with this product. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Area Control of Ticks: Use Dursban 2 EC to control ticks infesting non-crop land areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply Dursban 2 EC in water at the rate of 1 pint per acre (equivalent to 1 fluid ounce per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.

Warning (Area Control of Ticks): Do not allow public use of treated areas during application or until spray has dried. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff into lakes, streams, ponds or other bodies of water is likely to occur. Fish and crustaceans may be killed if their waters are contaminated with this product.

Mosquitoes: Apply Dursban 2 EC to mosquito infested, non-crop areas according to the conditions and rates shown in the following table. Mix Dursban 2 EC with either water or oil, such as kerosene or No. 2 fuel oil, using sufficient spray volume to give uniform coverage of area treated. Apply the spray using suitable application equipment such as hand and power ground sprayers, mist applicators, and aerial spray equipment.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold only in this unbroken package.

CONDITIONS and RATES to USE for MOSQUITO

Condition of Area to be Treated	Amount to Use per Acre	
	Dursban 2 EC (fluid ounces)	Chlorpyrifos (Pounds)
Larval Control		
None to medium vegetative cover	0.8 to 1.6	0.0125 to 0.025
Medium to heavy vegetative cover	1.6 to 3.2*	0.025 to 0.05*
Adult Control		
Light to medium vegetative cover	1.6	0.025
Medium to heavy vegetative cover	3.2*	0.05*

*The higher dosage will also give faster kill and longer residual control where there is no vegetative cover.

Warning (Mosquito Control): Dursban 2 EC is toxic to shrimp, crabs and fish. Do not apply to water containing crustaceans and fish. Consult your State Game Department before using this product for mosquito control.

Agway Inc.
Fertilizer-Chemical Division
Box 1333
Syracuse, New York 13201

EPA Reg. No. 8590-394

EPA Est. 8590-PA-1