WEBFOOT WEED AND FEED

WEBFOOT WEED AND FEED is a combination Plant Food and Selective Weedkiller. It supplies a balanced combination of Plant Food Nutrients to maintain a green, healthy lawn; Contains the Herbicide 2,4-D to kill the broadleaved weeds, Plantain and Dandelions, and the Herbicide Silvex to kill clovers and chickweed.

DIRECTIONS

SPREAD CONTENTS OF THIS BAG EVENLY OVER 2,500 sq. ft of LAWN. (50 x 50) Set spreader at a low serting or so that the hole openings are slightly smaller than 1/4 inch in diameter. See spreader settings bolow.


- 1. Apply when weeds are present and growing vigorously, preterably when they are young.
- 2. Application should be made following light rain, dew or light sprinkling previous to applying. (Grass should be dump enough so that WEBFOOT WEED AND FEED will adhere.)
- 3. Apply on a windstill day. The temperature should be at least 60° but not above 90° for 2 days following application.
- 4. Apply 3 days after mowing when weeds are actively growing.
- 5. Do not water treated lawn for 2 days or mow for 5 days after treatment.
- 6. Do not apply to new lawns until they are well established.

NOTE: Heavy rainfall, watering or traffic on treated area within 24 hours after application will minimize results.

Some browning of grass may appear after application but will normally disappear following next mowing.


GUARANTEED PLANT FOOD ANALYSIS

		Available Potash (Soluble K_2O) 4	19
Available Phosphoric Acid (P2Os) .	4%	fron (Fe)	9
		Sulphur (Combined) 4	19





Feeds 4









NOTE: Heavy rainfall, watering or traffic on treated area within 24 hours after application will minimize results.

> Some browning of grass may appear after application but will normally disappear following next mowing.

GUARANTEED PLANT FOOD ANALYSIS

Total Nitrogen	8%	Available Potash (Soluble K ₂ 0)	4%
Available Phosphoric Acid (P2O5).	4% 11		
		Sulphur (Combined)	4%

!NGREDIENTS

Sulphate of Ammonia, Amnonium Phosphates, Ferrous Sulphate, Muriate of Potash, Natural Gypsum, Tankage, 2,4-D-Silvex 30-10 Granular.

Spreaden Setting"

,		Lawn Beauty 16"	Central	Scott
	20 lbs. per 2500 Sq. ft.	No. 6	No. 8	No. 8

Note - Do NOT spread with Cyclone type speader.

CAUTIONS

Use of this product may be injurious to Bentgrass, St. Augustine Grass, Dichendra, Carpetgrass, or Clover Lawns.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

FREIGHT SHIPPING BAG


Meeting requirements of APPLICABLE FREIGHT CLASSIFICATION Sucrenteed by

BEIMS COMPANY, INC.


VARICCIOVER, WASH

ANTI-SKID 3-137-1/2














ACTIVE INGREDIENTS: 2, 4-Dichlorophend Isooctyl ester 2-(2, 4, 5-Trichloro

INERT INGREDIENTS:

WEBFCOT FERT

KEEP OUT

Inches comiting if swallowed. Avaid contact with skin. Wash should inapplied carefully on a day with low wind veloc insectables, fungicides, seeds or fertilizer. Rinse fertilizer contamination of feed and foodstuffs.

EPA

^{*}Spreader settings are suggestions only; condition of spreader, turf condition, and speed of walk while spreading may make a difference.