ACCEPTED

LAWNS LAWNS

\$6.95


DERAL



• • ; ; • •

MANUAL PROPERTY.





Siduron INERT INGR

*TUPERSA

LAWN LIFE CRAE and hairy crabgr other annual wea damage most of

Apply to the law tained when app gested that applications

NEWLY SEEDED

1. Apply conter

gested that application be avoided during February or early March.

NEWLY SEEDED LAWNS

- 1. Apply contents evenly over 5,000 square feet immediately after seeding (1.6 lbs. per 1,000 sq. ft.)
- 2. Provide sufficient moisture to insure germination of new grass seed.
- 3. An application of Lawn Life Turf Food is recommended after new turf is growing well.

ESTABLISHED, OVERSEEDED OR PATCHED LAWNS

- 1. Apply contents evenly over 2,500 square feet before crabgrass germinates (31/4 lbs. per 1,000 sq. 11.)
- 2. For best results, the lawn should be watered thoroughly soon after application.
- 3. Application of Lawn Life Turf Food is recommended in the Summer and Fall.

SUGGESTED SPREADER SETTINGS

	<u>2500 Sq. Ft.</u>	5000 Sq. Ft.
Lawn Beauty	7	5
Scott's	6	4
Jackson	5	3

These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

CAUTION: Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash hands with soap and water after handling or using. May irritate eyes, nose and skin. Do not apply to flowers, vegetables and ornamentals. Avoid contamination of foodstuff, ornamental fish ponds, irrigation ditches or water used for domestic purposes. Apply this product only as specified on this label.

IMPORTANT: Do not use on other than Penncross, Seaside, Highland and Astoria bentgrass strains nor on Bermuda grass as injury may result.

FEDERAL CHEMICAL COMPANY LOUISVILLE, KENTUCKY 40202

F-12-45A

FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION

Guaranteed by W-102

NP-6265

1167